arXiv:cond-mat/0312519v1 [cond-mat.str-el] 19 Dec 2003

The 2-matrix of the spin-polarized electron gas: contraction sum rules and spectral resolutions

P. Ziesche
Max-Planck-Institut für Physik komplexer Systeme,
Nöthnitzer Str. 38, D-01187 Dresden, Germany
and
F. Tasnádi
Leibniz-Institut für Festkörper- und Werkstoffforschung Dresden, Germany,
and University of Debrecen, Hungary

Keywords: spin-polarized electron gas, density matrices, cumulant expansion, sum rules, geminals

PACS numbers: 71.10.Ca, 05.30.Fk, 71.45.Gm

Abstract

The spin-polarized homogeneous electron gas with densities $\rho_\uparrow$ and $\rho_\downarrow$ for electrons with spin 'up' ($\uparrow$) and spin 'down' ($\downarrow$), respectively, is systematically analyzed with respect to its lowest-order reduced densities and density matrices and their mutual relations. The three 2-body reduced density matrices $\gamma_{\uparrow\uparrow}$, $\gamma_{\downarrow\downarrow}$, γ_a are 4-point functions for electron pairs with spins $\uparrow\uparrow$, $\downarrow\downarrow$, and antiparallel, respectively. From them, three functions $G_{\uparrow\uparrow}(x,y)$, $G_{\downarrow\downarrow}(x,y)$, $G_a(x,y)$, depending on only two variables, are derived. These functions contain not only the pair densities according to $g_{\uparrow\uparrow}(r) = G_{\uparrow\uparrow}(0,r)$, $g_{\downarrow\downarrow}(r) = G_{\downarrow\downarrow}(0,r)$, $g_a(r) = G_a(0,r)$ with $r = |\mathbf{r}_1 - \mathbf{r}_2|$, but also the 1-body reduced density matrices $\gamma_\uparrow$ and $\gamma_\downarrow$ being 2-point functions according to $\gamma_s = \rho_s f_s$ and $f_s(r) = G_{ss}(r,\infty)$ with $s = \uparrow, \downarrow$ and $r = |\mathbf{r}_1 - \mathbf{r}_1'|$. The contraction properties of the 2-body reduced density matrices lead to three sum rules to be obeyed by the three key functions G_{ss}, G_a. These contraction sum rules contain corresponding normalization sum rules as special cases. The momentum distributions $n_\uparrow(k)$ and $n_\downarrow(k)$, following from $f_\uparrow(r)$ and $f_\downarrow(r)$ by Fourier transform, are correctly normalized through $f_s(0) = 1$. In addition to the non-negativity conditions $n_s(k), g_{ss}(r), g_a(r) \geq 0$

[these quantities are probabilities], it holds $n_s(k) \leq 1$ and $g_{ss}(0) = 0$ due to the Pauli principle and $g_a(0) \leq 1$ due to the Coulomb repulsion. Recent parametrizations of the pair densities of the spin-unpolarized homogeneous electron gas in terms of 2-body wave functions (geminals) and corresponding occupancies are generalized (i) to the spin-polarized case and (ii) to the 2-body reduced density matrix giving thus its spectral resolutions.

1. Introduction

The homogeneous electron gas (HEG) is an important and widely used model for the phenomenon called electron correlation [1, 2, 3]. Its model parameters are $\rho_\uparrow$ and $\rho_\downarrow$, the densities of electrons with spin-up ($\uparrow$) and spin-down ($\downarrow$), respectively. Equivalent parameters are the density $\rho = \rho_\uparrow + \rho_\downarrow$ (from which follows the density parameter r_s defined by $4\pi r_s^3/3 = 1/\rho$ in a.u.) and $\zeta = (\rho_\uparrow - \rho_\downarrow)/\rho$, the spin-polarization. r_s is the (Wigner) radius of a sphere containing on average one electron. It measures simultaneously the coupling or interaction strength. The spin-dependent Wigner radii $r_{s\uparrow}$ and $r_{s\downarrow}$ are analogously defined by $4\pi r_{s\uparrow}^3/3 = 1/\rho_\uparrow$ and $4\pi r_{s\downarrow}^3/3 = 1/\rho_\downarrow$, respectively. This (jellium) model is relevant for the understanding of many effects in simple metals and semiconductors and it plays a crucial role in providing input quantities for approximate approaches to the many-electron problem of nonuniform density in solid-state theory and quantum chemistry. So the local-spin density approximation (LSD) of the density functional theory relies on $e(r_s, \zeta)$, the energy per particle of the HEG. Corresponding LSD (and beyond-LSD) functionals $E_{\text{xc}}[\rho_\uparrow, \rho_\downarrow]$ often treat the electron-electron interaction as a whole, without splitting it into its spin-resolved ($\uparrow\uparrow$, $\downarrow\downarrow$, and 'a' for spin-antiparallel) components. But some of the most popular functionals are constructed by considering spin-resolved contributions to the correlation energy. This makes the recent interest in the spin-polarized HEG understandable. The system under consideration has the two afore mentioned model parameters r_s and ζ. Its ground state is characterized by the reduced densities $n_\uparrow(k)$, $n_\downarrow(k)$ [the momentum distributions] and $g_{\uparrow\uparrow}(r)$, $g_{\downarrow\downarrow}(r)$, $g_a(r)$ [the pair densities (PDs)]. Corresponding energies are $t_\uparrow$, $t_\downarrow$ [the kinetic energies] and $v_{\uparrow\uparrow}$, $v_{\downarrow\downarrow}$, v_a [the interaction energies] such that the total bulk energy e is given by $e = e_\uparrow + e_\downarrow$ where $e_\uparrow = t_\uparrow + v_\uparrow$ and $e_\downarrow = t_\downarrow + v_\downarrow$ with $v_\uparrow = v_{\uparrow\uparrow} + v_a$ and $v_\downarrow = v_{\downarrow\downarrow} + v_a$. All these quantities depend parametrically on r_s and ζ. The PDs have been studied recently in Refs. [4, 5, 6, 7, 8]. Recent studies of $n(k)$ for the spin-unpolarized case are in Ref. [9]. The total (spin-weighted) PD $g = (\rho_\uparrow/\rho)^2 g_{\uparrow\uparrow} + (\rho_\downarrow/\rho)^2 g_{\downarrow\downarrow} + 2(\rho_\uparrow\rho_\downarrow/\rho^2) g_a$ is related to the total momentum distribution $n(k) = \frac{1}{2}[n_\uparrow(k) + n_\downarrow(k)]$ (i) by the virial theorem [10, 11] and (ii) through the 2-body reduced density matrix (2-matrix) [12, 13, 14, 15]. The latter means: its diagonal elements give the PDs and its contraction yields the 1-matrix, which is the Fourier transform of the momentum distribution.

These mutual relations between the PDs and the momentum distributions are investigated for the spin-polarized HEG in the following. In detail its 2-matrix and quantities derived from it are systematically analyzed in particular in terms of the already mentioned contraction sum rules (SRs) and the so-called spectral resolutions. This more general analysis (with introductory parts in Secs. 2-6 and the main parts in Secs. 7-9) includes the special cases of 'no interaction' (but $\zeta \neq 0$), 'no polarization' (but $r_s \neq 0$), and 'full polarization' (with $\zeta = 1$ and $r_s \neq 0$) [they are summarized in Apps. 1, 2, 3, respectively]. Thereby four questions are considered:

1. For the spin-unpolarized HEG it is well-known as already mentioned, that the combined virial and Hellmann-Feynman theorem relates the interaction energy v to the kinetic energy t, such that $t(r_s)$ can be calculated provided $v(r_s)$ is known from the PD $g(r; r_s)$. The question arises whether there are theorems for the spin-polarized HEG, such that not only t, but also its spin-components $t_\uparrow$ and $t_\downarrow$ can be calculated from $v_{\uparrow\uparrow}$, $v_{\downarrow\downarrow}$, v_a, which are available because $g_{\uparrow\uparrow}$, $g_{\downarrow\downarrow}$, g_a are known as functions of r and r_s, ζ [4, 5, 6, 7, 8]. The answer in Sec. 3 is 'no' (disagreeing with a formula in Ref. [16]).
2. The virial theorem is an integral relation between $n(k)$ and $g(r)$. Other relations arise from their common origin, namely the 2-body reduced density matrix. The question 'how these relations simplify themselves for homogeneous systems ?' is answered in Sec. 8.
3. Kimball [17] and Overhauser [18] introduced a parametrization of the HEG-PD in terms of 2-body wave functions (so-called geminals), which are the solutions of a 2-body Schrödinger equation with an appropriately chosen effective interaction (screened Coulomb repulsion), which describes the scattering of two electrons in the medium of all the others. Further developments of this idea [19, 20] allows one to calculate the PD in excellent agreement with the results of quantum Monte Carlo calculations [4, 6]. Because nowadays the PDs not only for $\zeta = 0$ but also for $\zeta \neq 0$ are available [6, 7, 8], the question may be posed whether in this more general spin-polarized case the PDs can be parametrized again in terms of geminals following from a screened Coulomb repulsion. In Sec. 9 the answer

'yes' is assumed and a corresponding analysis is presented as the necessary pre-stage of a numerical study.
4. In Ref. [21] it has been shown how the PD-normalization can be expressed in terms of scattering phase shifts of the geminals similar as this is known from the Friedel sum rule. Moreover in Ref. [22] the Kimball-Overhauser approach is generalized with the assumption that the PD-geminals can be used to represent not only the PD but also the 2-matrix. The contraction of which gives the 1-matrix and thus the momentum distribution. Along this line the normalization SR of Ref. [21] is generalized in Ref. [22] to a contraction SR, which allows one in principle to self-consistently calculate the momentum distribution provided the afore mentioned geminals are known. For a summary cf. Ref. [23]. Again the question arises whether this analysis can be extended to non-vanishing spin-polarization. Preliminary considerations are presented in Sec. 10.

The paper is structured as it follows. Sec. 1 defines the HEG by its Hamiltonian and lists the ground state quantities. In Sec. 2 the virial and Hellmann-Feynmam theorems are revisited. Secs. 4 and 5 present the 1-matrix γ_1 and the 2-matrix γ_2, respectively, with the spin-structure of γ_2, derived in App. 4. Sec. 6 summarizes the four PDs and their normalizations. Sec. 7 describes how the 1-matrix can be obtained from the 2-matrix by contraction. Apps. 1, 2, 3 summarize the special cases of 'no interaction', 'no spin-polarization', 'full spin-polarization', respectively. In Sec. 8 the 2-matrix is discussed in terms of Wick's theorem and size-extensivity, guaranteed by linked diagrams and cumulants. Sec. 9 presents the spectral resolutions of the 2-matrix and the cumulant 2-matrix in terms of geminals and corresponding occupancies. Sec. 10 expresses the normalization and contraction SRs in terms of the scattering phase shifts which describe the asymptotics of the geminals.

2. The system and its ground state properties

The GS of the spin-polarized HEG is characterized by the average densities $\rho_\uparrow$ and $\rho_\downarrow$ of the spin-up, respectively spin-down electrons or equivalently by the average spin-summed density $\rho = \rho_\uparrow + \rho_\downarrow$ and the spin-polarization $\zeta = (\rho_\uparrow - \rho_\downarrow)/\rho$, thus $\rho_\uparrow/\rho = (1+\zeta)/2$ and $\rho_\downarrow/\rho = (1-\zeta)/2$. Essential parts of the phenomenon 'electron correlation' are contained in the two dimensionless momentum distributions $n_\uparrow(k)$, $n_\downarrow(k)$ and in four dimensionless PDs, namely $g_{\uparrow\uparrow}(r)$, $g_{\downarrow\downarrow}(r)$, and $g_\pm(r)$ or equivalently $g_a = (g_+ + g_-)/2$, $\Delta g = (g_+ - g_-)/2$. The index '$\pm$' stands for singlet/triplet, 'a' for spin-antiparallel, respectively. To get these GS quantities one has to solve

$$\hat{H}\Psi = E\Psi, \quad \hat{N}_s\Psi = N_s\Psi \tag{1}$$

with $\hat{H} = \hat{T} + \hat{V}$ and

$$\begin{aligned} \hat{T} &= \int d1\, \hat{\psi}_1^\dagger t_1 \hat{\psi}_1 \ , \qquad t_1 = -\frac{\hbar^2}{2m}\left(\frac{\partial}{\partial \mathbf{r}_1}\right)^2 , \\ \hat{V} &= \int \frac{d1d2}{2!}\, [:(\hat{\rho}_1 - \rho_{\Omega 1})v_{12}(\hat{\rho}_2 - \rho_{\Omega 2}):] \ , \qquad v_{12} = \frac{\epsilon^2}{r_{12}}, \quad r_{12} = |\mathbf{r}_1 - \mathbf{r}_2|, \\ \hat{\rho}_1 &= \hat{\psi}_1^\dagger \hat{\psi}_1, \quad :\hat{\rho}_1\hat{\rho}_2 := \hat{\psi}_1^\dagger \hat{\psi}_2^\dagger \hat{\psi}_2 \hat{\psi}_1 \ , \qquad \rho_{\Omega 1} = \rho_{s_1}\Theta_\Omega(\mathbf{r}_1) \end{aligned} \tag{2}$$

and to take the TDL $N_s, \Omega \to \infty$, $N_s/\Omega = \text{const} = \rho_s$ with $N = N_\uparrow + N_\downarrow$. Here $1 = (\mathbf{r}_1, \sigma_1)$, σ and $s = \uparrow$ or $\downarrow$, and $\hat{\psi}_1 = \hat{\psi}_{s_1}(\mathbf{r}_1)$. $\hat{\psi}_s^\dagger(\mathbf{r})$ and $\hat{\psi}_s(\mathbf{r})$ are the Fermion creation and annihilation field operators such that $\hat{\rho}_s(\mathbf{r}) = \hat{\psi}_s^\dagger(\mathbf{r})\hat{\psi}_s(\mathbf{r})$ is the density operator for electrons with spin s and $\hat{\rho}(\mathbf{r}) = \hat{\rho}_\uparrow(\mathbf{r}) + \hat{\rho}_\downarrow(\mathbf{r})$ is the spin-summed density operator. Hence, $\hat{N}_s = \int d^3r\, \hat{\psi}_s^\dagger(\mathbf{r})\hat{\psi}_s(\mathbf{r})$ and $\hat{N} = \hat{N}_\uparrow + \hat{N}_\downarrow$ are the particle-number operators for spin s and spin-summed, respectively. The operators $(:\hat{\rho}_1\hat{\rho}_2:)$, $\hat{\rho}_1(\rho_\Omega)_2 + (\rho_\Omega)_1\hat{\rho}_2$, and $(\rho_\Omega)_1(\rho_\Omega)_2$ describe the electron-electron $(--)$, electron-background $(-+$ and $+-)$, and background-background $(++)$ interaction, respectively. Corresponding expectation values are the spin-density $\rho_s(\mathbf{r}) = \langle\hat{\rho}_s(\mathbf{r})\rangle$ for $s = \uparrow, \downarrow$, as well as the two 1-matrices $\rho_s f_s(|\mathbf{r}_1 - \mathbf{r}_1'|) = \langle\psi_s^\dagger(\mathbf{r}_1')\psi_s(\mathbf{r}_1)\rangle$ and the four dimensionless PDs $g_{ss}, g_a, \Delta g$ defined by $\rho_s^2 g_{ss}(r_{12}) = \langle:\hat{\rho}_s(\mathbf{r}_1)\hat{\rho}_s(\mathbf{r}_2):\rangle$, $\quad \rho_\uparrow\rho_\downarrow g_a(r_{12}) = \langle\hat{\rho}_\uparrow(\mathbf{r}_1)\hat{\rho}_\downarrow(\mathbf{r}_2)\rangle$, and $\rho_\uparrow\rho_\downarrow\Delta g(r_{12}) = \langle\hat{\psi}_\downarrow^\dagger(\mathbf{r}_1)\hat{\psi}_\uparrow^\dagger(\mathbf{r}_2)\hat{\psi}_\downarrow(\mathbf{r}_2\hat{\psi}_\downarrow(\mathbf{r}_1)\rangle$. Then the spin-weighted PD $g(r)$ is defined by $\rho^2 g(r) = \rho_\uparrow^2 g_{\uparrow\uparrow}(r) + \rho_\downarrow^2 g_{\downarrow\downarrow}(r) + 2\rho_\uparrow\rho_\downarrow g_a(r)$. The momentum distributions are given by $n_s(k) = \langle\hat{a}_{s\mathbf{k}}^\dagger\hat{a}_{s\mathbf{k}}\rangle$ and $n(k) = \frac{1}{2}[n_\uparrow(k) + n_\downarrow(k)]$ with $\sum_\mathbf{k} n_s(k) = N_s$ and $2\sum_\mathbf{k} n(k) = N$; the normalizations per particle are $\frac{1}{N}\sum_\mathbf{k} n_\uparrow(k) = \frac{1}{2}(1+\zeta) > \frac{1}{2}$, $\frac{1}{N}\sum_\mathbf{k} n_\downarrow(k) = \frac{1}{2}(1-\zeta) < \frac{1}{2}$, and $\frac{2}{N}\sum_\mathbf{k} n(k) = 1$.

3. The virial and Hellmann-Feynman theorems

From the five mentioned reduced densities $n_\uparrow(k), n_\downarrow(k), g_{\uparrow\uparrow}(r), g_{\downarrow\downarrow}(r), g_a(r)$ follow five contributions to the total energy $E = T + V$ with $T = T_\uparrow + T_\downarrow$ and $V = V_{\uparrow\uparrow} + V_{\downarrow\downarrow} + 2V_a$, being the GS expectation values of $\hat{H} = \hat{T} + \hat{V}$ with $\hat{T} = \hat{T}_\uparrow + \hat{T}_\downarrow$ and $\hat{V} = \hat{V}_{\uparrow\uparrow} + \hat{V}_{\downarrow\downarrow} + 2\hat{V}_a$. Now, the virial theorem is one relation between T and V with the consequence: if V is known, then T can be calculated. The question arises: Are there perhaps two virial theorems, one for spin $\uparrow$ and one for spin $\downarrow$, which - for given PDs g_{ss} and g_a - would possibly allow one to calculate the spin components $T_\uparrow$ and $T_\downarrow$, separately ? The answer is ‘no’. To show this, one may define $\hat{H}_s = \hat{T}_s + \hat{V}_s$, $\hat{V}_s = \hat{V}_{ss} + \hat{V}_a$ and $E_s = T_s + V_s$, $V_s = V_{ss} + V_a$, for $s = \uparrow, \downarrow$, such that $\hat{H} = \hat{H}_\uparrow + \hat{H}_\downarrow$ and $E = E_\uparrow + E_\downarrow$, but there is only one virial theorem, to be derived from the hypervirial theorem $\langle[\hat{F}, \hat{H}]\rangle = 0$, which holds for an eigenstate Ψ of $\hat{H}$ and an arbitrary operator $\hat{F}$ of the system; the commutator is denoted by $[A, B] = AB - BA$. With the virial operator $\hat{F} = \hat{\mathbf{rp}}$ and with its spin resolution $\hat{\mathbf{rp}} = (\hat{\mathbf{rp}})_\uparrow + (\hat{\mathbf{rp}})_\downarrow$, $(\hat{\mathbf{rp}})_s = \int d^3r\, \hat{\psi}_s^\dagger(\mathbf{r})\mathbf{rp}\hat{\psi}_s(\mathbf{r})$ it is easy to show

$$\begin{aligned} 0 = -\frac{\mathrm{i}}{\hbar}\langle[\hat{\mathbf{rp}}, \hat{H}]\rangle &= \langle[\widehat{\mathbf{p}\frac{\partial}{\partial\mathbf{p}}}, \hat{T}]\rangle + \langle[(\widehat{-\mathbf{r}\frac{\partial}{\partial\mathbf{r}}}), \hat{V}]\rangle \\ &= 2T + V + 3\Omega\langle\frac{\partial\hat{V}}{\partial\Omega}\rangle, \quad \frac{\partial\hat{V}}{\partial\Omega} = \frac{\partial\hat{H}}{\partial\Omega}. \end{aligned} \tag{3}$$

Only $\hat{V}^{+-} + \hat{V}^{-+} + V^{++}$ depend on the system parameter Ω = volume. And there is only one Hellmann-Feynman theorem [24]

$$\frac{\partial E}{\partial\Omega} = \langle\frac{\partial\hat{H}}{\partial\Omega}\rangle. \tag{4}$$

The reason: There is only one MB Schrödinger equation $(\hat{H} - E)\Psi = 0$. In addition to the virial theorem (3) one may derive the relations, cf. [25]

$$0 = \langle[(\hat{\mathbf{rp}})_\uparrow, \hat{H}]\rangle = 2T_\uparrow + V_{\uparrow\uparrow} + X + 3\Omega\langle\frac{\partial}{\partial\Omega}\hat{H}_\uparrow\rangle, \tag{5}$$

$$0 = \langle[(\hat{\mathbf{rp}})_\downarrow, \hat{H}]\rangle = 2T_\downarrow + V_{\downarrow\downarrow} - X + 3\Omega\langle\frac{\partial}{\partial\Omega}\hat{H}_\downarrow\rangle, \tag{6}$$

where the expection value of

$$\hat{X} = \frac{1}{2}\int d^3r_1 d^3r_2\, \hat{\rho}_\uparrow(\mathbf{r}_1)\hat{\rho}_\downarrow(\mathbf{r}_2)\frac{r_1^2 - r_2^2}{r_{12}^2}\frac{\epsilon^2}{r_{12}}. \tag{7}$$

appears. In Ref. [25] the modified (‘filtered’ with the prefactor $\mathbf{Rr}/r^2 = R\xi/r$, where $\mathbf{R}$ is the centre-of-mass and $\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2$) Coulomb repulsion in Eq.(7) is referred to as ‘spin polarization potential’. Note $\langle\partial\hat{H}_s/\partial\Omega\rangle \neq \partial E_s/\partial\Omega$: The HF theorem (4) does not split into two different theorems for spin $\uparrow$ and spin $\downarrow$. So, the sum of Eqs.(5) and (6) yields correctly the VT (3), but their difference gives

$$2\Delta T + \Delta V + 2X + 3\Omega\langle\frac{\partial}{\partial\Omega}\Delta\hat{H}\rangle = 0 \tag{8}$$

with $\Delta T = T_\uparrow - T_\downarrow$, $\Delta V = V_{\uparrow\uparrow} - V_{\downarrow\downarrow}$ (V_a cancels), $\Delta\hat{H} = \hat{H}_\uparrow - \hat{H}_\downarrow = \Delta\hat{T} + \Delta\hat{V}$.

Specification of Ω: For an atom with a pointlike positive background charge, i.e. $\Omega \to 0$, $\rho \to \infty$, $\rho\Omega \to Z$ (atomic limit), all the terms with $\partial/\partial\Omega$ are no longer present. Then the virial theorem (3) is simply $2T + V = 0$ and the other relation (8) is $2\Delta T + \Delta V + 2X = 0$, in Ref. [25] called 'spin virial theorem'. Contrary to this for the spin-polarized HEG in its TDL with $N, \Omega \to \infty$ and $t = T/N, v = V/N, e = E/N$, and $3\Omega\partial e/\partial\Omega = r_s\partial e/\partial r_s$, from the theorems (3) and (4) follows the bulk virial theorem [10]

$$2t + v + r_s\frac{\partial}{\partial r_s}e = 0 \quad \text{or} \quad \frac{1}{r_s}\frac{\partial}{\partial r_s}r_s^2 t + \frac{\partial}{\partial r_s}r_s v = 0, \tag{9}$$

where the derivative is related to the pressure of the system: $r_s\partial e/\partial r_s = -3p/\rho$. For the case 'no interaction' the virial theorem (9) simply says $t^0 \sim 1/r_s^2$. Also the surface virial theorem $2t_{\rm surf} + v_{\rm surf} + (r_s\partial/\partial r_s + 2)e_{\rm surf} = 0$ can be derived [26]. But what results from Eq.(8)? One may write the sum and the difference of the last terms on the rhs's of Eqs.(5) and (6) as

$$3\Omega\langle\frac{\partial}{\partial\Omega}\hat{H}\rangle = 3\Omega\frac{\partial}{\partial\Omega}E - 3\Omega\left(\frac{\partial}{\partial\Omega}\right)_\Psi\langle\hat{H}\rangle \tag{10}$$

and

$$3\Omega\langle\frac{\partial}{\partial\Omega}\Delta\hat{H}\rangle = 3\Omega\frac{\partial}{\partial\Omega}\Delta E - 3\Omega\left(\frac{\partial}{\partial\Omega}\right)_\Psi\langle\Delta\hat{H}\rangle, \tag{11}$$

respectively. $(\partial/\partial\Omega)_\Psi$ means differentiation with respect to Ψ, not $\hat{H}$. Now, the last term on the rhs of Eq.(10) vanishes, because of the Hellmann-Feynman theorem (4), thus giving the bulk virial theorem (9). Contrary to this the last term on the rhs of Eq.(11) gives an expression, which does not vanish. So, Eq.(11) used in Eq.(8) gives only the trivial identity $0 = 0$. This is seen from

$$2\Delta t + \Delta v + r_s\frac{\partial\Delta e}{\partial r_s} = \frac{1}{r_s}\frac{\partial}{\partial r_s}r_s^2\Delta t + \frac{\partial}{\partial r_s}r_s\Delta v = r_s\left(\frac{\partial}{\partial r_s}\right)_\Psi\langle\frac{\Delta\hat{H}}{N}\rangle \tag{12}$$

and (in the TDL)

$$x = \frac{X}{N} = \frac{\rho_\uparrow\rho_\downarrow}{\rho^2}2\int d^3r\ \rho\int\frac{d^3R}{\Omega}\frac{R}{r}\int_{-1}^{+1}\frac{d\xi}{2}\xi g_\Omega(R, r, \xi)\frac{\epsilon^2}{r} = O(N^{-2/3}), \tag{13}$$

where $\rho_\uparrow\rho_\downarrow g_\Omega(R, r, \xi) = \langle\hat{\rho}_\uparrow(\mathbf{r}_1)\hat{\rho}_\downarrow(\mathbf{r}_2)\rangle$, $\mathbf{R} = (\mathbf{r}_1 + \mathbf{r}_2)/2$, $\mathbf{r} = (\mathbf{r}_1 - \mathbf{r}_2)$, $\xi = \sphericalangle(\mathbf{R}, \mathbf{r})$, and $g_\infty(R, r, \xi) = g_a(r)$. So, from the HEG-PDs $g_{\uparrow\uparrow}$, $g_{\downarrow\downarrow}$, g_a as functions of r and the parameters r_s, ζ one can calculate only t as a function of r_s and ζ, but not its spin components $t_\uparrow, t_\downarrow$, separately. Thus Eq.(3.11) of Ref. [16] is wrong. But one may define a quantity y as a function of r_s and ζ by

$$2\Delta t + \Delta v + r_s \frac{\partial \Delta e}{\partial r_s} = y \tag{14}$$

(so this is not a theorem !). For 'no interaction' it is $\Delta v^0 = 0, \Delta t^0 \sim 1/r_s^2$, so y vanishes with $r_s \to 0$. - In the literature there are hints on a magnetic virial theorem if one adds to the Hamiltonian (2) a Zeeman term $\sim \hat{S}_z = \frac{\hbar}{2}(\hat{N}_\uparrow - \hat{N}_\downarrow)$ [16, 27].

4. The 1-matrices

The 1-matrix is defined by $\gamma_1(1|1') = \langle \hat{\psi}^+_{1'} \hat{\psi}_1 \rangle$, normalized as $\mathrm{Tr}\gamma_1 = N_\uparrow + N_\downarrow = N$. In terms of natural spin-orbitals $\psi_k(1)$ and corresponding occupation numbers ν_k the spectral resolution of γ_1 is $\gamma_1(1|1') = \sum_k \psi_k(1)\nu_k\psi_k^*(1')$ with $0 < \nu_k < 1$ and $\sum_k \nu_k = N$. For a spin-independent Hamiltonian as $\hat{H}$ of Eqs.(1) and (2) it is $\psi_k(1) = \delta_{s,\sigma_1}\varphi_{\mathbf{k}}(\mathbf{r}_1)$ with $\langle \psi^\dagger_\uparrow(\mathbf{r}')\psi_\downarrow(\mathbf{r})\rangle = 0$ and $\langle \psi^\dagger_\downarrow(\mathbf{r}')\psi_\uparrow(\mathbf{r})\rangle = 0$. Thus

$$\gamma_1(1|1') = \delta_{\sigma_1,\sigma_1'}[\delta_{\sigma_1,\uparrow}\gamma_\uparrow(\mathbf{r}_1|\mathbf{r}_1') + \delta_{\sigma_1,\downarrow}\gamma_\downarrow(\mathbf{r}_1|\mathbf{r}_1')] \tag{15}$$

with

$$\gamma_s(\mathbf{r}|\mathbf{r}') = \sum_{\mathbf{k}} \varphi_{\mathbf{k}}(\mathbf{r})n_s(\mathbf{k})\varphi^*_{\mathbf{k}}(\mathbf{r}'), \mathrm{Tr}\gamma_s = \sum_{\mathbf{k}} n_s(\mathbf{k}) = N_s, \sum_{\mathbf{k}} = \int \frac{\Omega d^3k}{(2\pi)^3}. \tag{16}$$

The spin-traced 1-matrix is $\gamma = \gamma_\uparrow + \gamma_\downarrow = \rho_\uparrow f_\uparrow(r) + \rho_\downarrow f_\downarrow(r)$.

For a homogeneous system the natural orbitals are plane waves $\varphi_{\mathbf{k}}(\mathbf{r}) = \frac{1}{\sqrt{\Omega}}\mathrm{e}^{\mathrm{i}\mathbf{kr}}$ and it is $\gamma_s(\mathbf{r}|\mathbf{r}') = \rho_s f_s(|\mathbf{r}-\mathbf{r}'|)$, which defines the dimensionless 1-matrices

$$f_s(r) = \frac{1}{N_s}\sum_{\mathbf{k}} n_s(k)\mathrm{e}^{\mathrm{i}\mathbf{kr}}, \tag{17}$$

displayed in Fig. 1. The Fourier transforms of $f_s(r)$ give the momentum distributions $n_s(k)$ with $0 \le n_s(k) \le 1$ and with a large and a small Fermi sphere: $(k_{\mathrm{F}s})^3 = 6\pi^2\rho_s$, $k_{\mathrm{F}\uparrow} > k_{\mathrm{F}\downarrow}$. With $k_\mathrm{F}^3 = 3\pi^2\rho$ it is $k_{\mathrm{F}\uparrow}/k_\mathrm{F} = (1+\zeta)^{1/3}$ and $k_{\mathrm{F}\downarrow}/k_\mathrm{F} = (1-\zeta)^{1/3}$. $f_s(0) = 1$ makes the $n_s(k)$ correctly normalized: $\sum_{\mathbf{k}} n_s(k) = N_s$. The deviations of $n_s(k)$ from 0 and 1 (its non-idempotency) measure the correlation strength: $\sum_{\mathbf{k}}[n_s(k)]^2 < N_s$. The spin-summed momentum distribution is $n(k) = \frac{1}{2}[n_\uparrow(k) + n_\downarrow(k)]$. Whether the asymptotics of the $n_s(k)$ is spin-independent has to be checked. If so, it should hold $n_s(k \to \infty) = (2\alpha_0/3\pi)^2 g_a(0)/k^8$, $\alpha_0 = (4/9\pi)^{1/3}r_s$ [28, 29].

For the special cases 'no interaction', 'no polarization', and 'full polarization' cf. Apps. 1, 2, and 3, respectively.

The recently available momentum distribution $n(k, r_s)$ of the spin-unpolarized HEG (k measured in units of $k_{\rm F}$) [9] can be used to approximately construct the momentum distributions of the spin-polarized HEG for small values of ζ:

$$n_\uparrow(k, r_s, \zeta) \approx n\left(\frac{k}{(1+\zeta)^{1/3}}, \frac{r_s}{(1+\zeta)^{1/3}}\right), \quad n_\downarrow(k, r_s, \zeta) \approx n\left(\frac{k}{(1-\zeta)^{1/3}}, \frac{r_s}{(1-\zeta)^{1/3}}\right). \quad (18)$$

This allows one to calculate the above mentioned spin components $t_\uparrow(r_s, \zeta)$ and $t_\downarrow(r_s, \zeta)$ of the total kinetic energy $t(r_s, \zeta) = t_\uparrow(r_s, \zeta) + t_\downarrow(r_s, \zeta)$ and to control it by comparing with $t(r_s, \zeta)$, calculated from the PDs with the help of the virial theorem (9). Results for $\zeta = 1/3$ and $r_s < 10$ are displayed in Figs. 2 and 3. They show that the approximation (18) affords the correct $t(r_s, \zeta)$.

Related quantities derived from $f_\uparrow$, $f_\downarrow$ or $n_\uparrow$, $n_\downarrow$ are the 1-matrices (referred to as non-idempotency matrices)

$$\beta_s(r_{11'}) = \left(\frac{\rho_s}{\rho}\right)^2 \int d^3r_2 \rho f_s(r_{12}) f_s(r_{21'}) = \frac{1}{N}\sum_{\mathbf{k}} [n_s(k)]^2 {\rm e}^{i\mathbf{k}\mathbf{r}_{11'}} \quad , \qquad \beta_s(0) = b_s,$$

$$\beta_a(r_{11'}) = \frac{\rho_\uparrow \rho_\downarrow}{\rho^2} \int d^3r_2 \rho f_\uparrow(r_{12}) f_\downarrow(r_{21'}) = \frac{1}{N}\sum_{\mathbf{k}} n_\uparrow(k) n_\downarrow(k) {\rm e}^{i\mathbf{k}\mathbf{r}_{11'}} \quad , \qquad \beta_a(0) = b_a. \quad (19)$$

The product $n_\uparrow(k)n_\downarrow(k)$ makes the $\mathbf{k}$-summation to run over three different regions: The inner sphere $k < (1-\zeta)^{1/3}$, the shell $(1-\zeta)^{1/3} < k < (1+\zeta)^{1/3}$, and the outer region $k > (1+\zeta)^{1/3}$. The quantities b_s and b_a are related to spin-generalized Löwdin parameters

$$\begin{aligned} c_s = 1 - \frac{N}{N_s} b_s &= 1 - \frac{1}{N_s}\sum_{\mathbf{k}} [n_s(k)]^2, \\ c_a = 1 - \frac{N}{N_\downarrow} b_a &= 1 - \frac{1}{N_\downarrow}\sum_{\mathbf{k}} n_\uparrow(k) n_\downarrow(k), \end{aligned} \quad (20)$$

which vanish for 'no interaction', cf. App. 1. For 'no polarization' (cf. App. 2) it becomes $n_\uparrow(k) = n_\downarrow(k) \equiv n(k)$ and therefore $b_s = b_a \equiv b$, $c_s = c_a \equiv c$, $c = 1 - 2b$, and $b = \frac{1}{N}\sum_{\mathbf{k}}[n(k)]^2$. It was already P.-O. Löwdin who has asked for the meaning of ${\rm Tr}\gamma_1^2$. The answer is nowadays: $1 - {\rm Tr}\gamma_1^2/N$ is the normalization of the cumulant 2-matrix and one possible index of the correlation strength. As seen from $c = \frac{2}{N}\sum_{\mathbf{k}} n(k)[1-n(k)]$, this expression is particle-hole symmetric [31]. In addition to the above defined non-idempotency matrices, the 1-matrices $\alpha(r)$, $\tilde{\alpha}(r)$ appear, when contracting the 2-matrix, cf. Eq.(36). They cause (for spin-antiparallel electron pairs) the difference between singlet and triplet PDs and describe fluctuations, which mimick spin-flip processes. This means the following: One may consider a spatial part (region, domain, or fragment) ω of the system, say a sphere, which contains on average $\omega\rho$ electrons. One may furthermore ask for the probability of finding $N_\omega \neq \omega\rho$ electrons in ω. This can be realized by a certain number $N_{\omega\uparrow}$ of spin-up electrons and a certain number $N_{\omega\downarrow}$ of spin-down electrons. These numbers can change under the constraint $N_{\omega\uparrow} + N_{\omega\downarrow} = N_\omega$, what looks effectively as if electrons in ω had made spin-flips, although such fluctuations are based only on electrons (with spin-up and spin-down) leaving or entering the region ω. In general, for particle number fluctuations in spatial regions cf. Ref. [32]. - For 'no interaction' it is $\alpha^0(r) = \frac{N_\downarrow}{N} f_\downarrow^0(r)$ and $\tilde{\alpha}^0(r) = 0$, cf. App. 1.

5. The 2-matrices

The 2-matrix $\gamma_2(1|1', 2|2') = \langle \hat{\psi}_{1'}^\dagger \hat{\psi}_{2'}^\dagger \hat{\psi}_2 \hat{\psi}_1 \rangle$ is with $\hat{N}_{\sigma_2} \hat{\psi}_1 = \hat{\psi}_1 (\hat{N}_{\sigma_2} - \delta_{\sigma_1 \sigma_2})$, hence $\hat{N}\hat{\psi}_1 = \hat{\psi}_1(\hat{N} - 1)$, normalized as $\mathrm{Tr}\gamma_2 = N(N-1)$ with $\mathrm{Tr} \cdots = \int d1 d2 \cdots |_{1'=1, 2'=2}$. For given positions $\mathbf{r}_1, \mathbf{r}_1', \mathbf{r}_2, \mathbf{r}_2'$ there are 16 spin matrix elements. But only six of them are non-zero, the other 10 matrix elements vanish. With the short-hand notation $\langle \sigma_1' \sigma_2' \sigma_2 \sigma_1 \rangle = \langle \hat{\psi}_{\sigma_1'}^\dagger(\mathbf{r}_1') \hat{\psi}_{\sigma_2'}^\dagger(\mathbf{r}_2') \hat{\psi}_{\sigma_2}(\mathbf{r}_2) \hat{\psi}_{\sigma_1}(\mathbf{r}_1) \rangle$ the non-vanishing elements are

$$\begin{aligned}
\gamma_{\uparrow\uparrow} &= \langle \uparrow\uparrow\uparrow\uparrow \rangle, & \mathrm{Tr}\gamma_{\uparrow\uparrow} &= N_\uparrow(N_\uparrow - 1), \\
\gamma_{\downarrow\downarrow} &= \langle \downarrow\downarrow\downarrow\downarrow \rangle, & \mathrm{Tr}\gamma_{\downarrow\downarrow} &= N_\downarrow(N_\downarrow - 1), \\
\gamma_a^1 &= \langle \uparrow\downarrow\downarrow\uparrow \rangle, & \mathrm{Tr}\gamma_a^1 &= N_\downarrow N_\uparrow, \\
\gamma_a^2 &= \langle \downarrow\uparrow\uparrow\downarrow \rangle, & \mathrm{Tr}\gamma_a^2 &= N_\uparrow N_\downarrow, \\
\gamma_a^3 &= \langle \downarrow\uparrow\downarrow\uparrow \rangle, & \mathrm{Tr}\gamma_a^3 &= -Na, \\
\gamma_a^4 &= \langle \uparrow\downarrow\uparrow\downarrow \rangle, & \mathrm{Tr}\gamma_a^4 &= -Na,
\end{aligned} \tag{21}$$

whereas e.g. $\langle \uparrow\uparrow\downarrow\downarrow \rangle$, $\langle \uparrow\uparrow\uparrow\downarrow \rangle$ vanish. Each of these matrix elements depends on $\mathbf{r}_1|\mathbf{r}_1', \mathbf{r}_2|\mathbf{r}_2'$. The first two lines concern spin-parallel electron pairs and the next four lines describe spin-antiparallel electron pairs. The properties of these matrix elements (21) are listed in Table I together with other related matrix elements for the spin-antiparallel electrons and with the corresponding spin-matrices defined below in Eq.(22). The operators therein are defined as it follows: D creates the diagonal elements with $\mathbf{r}_1' = \mathbf{r}_1$ and $\mathbf{r}_2' = \mathbf{r}_2$, P makes the permutation $\mathbf{r}_1 \leftrightarrow \mathbf{r}_2$, Q effects the hermitian conjugation with $\mathbf{r}'_1 \leftrightarrow \mathbf{r}_1$, $\mathbf{r}'_2 \leftrightarrow \mathbf{r}_2$, and c.c. (= complex conjugation), and C causes the contraction with $\mathbf{r}'_2 \leftrightarrow \mathbf{r}_2$ and $\int d^3r_2$, where 1-matrices appear, namely $\gamma_\uparrow = \rho_\uparrow f_\uparrow$, $\gamma_\downarrow = \rho_\downarrow f_\downarrow$, and $\rho\alpha_3$, $\rho\alpha_4$. For the position matrices γ, Tr means $\mathbf{r}'_1 = \mathbf{r}_1$, $\mathbf{r}'_2 = \mathbf{r}_2$, and $\int d^3r_1 d^3r_2$. For the spin-matrices δ, Tr means $\sigma'_1 = \sigma_1$, $\sigma'_2 = \sigma_2$, and $\sum_{\sigma_{1,2}}$. Note $\gamma'_a = (\gamma_- - \gamma_+)/2$ and $\Delta g = (g_+ - g_-)/2$. Columns for $\tilde{\mathrm{D}}$ making $\mathbf{r}'_1 = \mathbf{r}_2$, $\mathbf{r}'_2 = \mathbf{r}_1$ and similarly for $\tilde{\mathrm{P}}$, $\tilde{\mathrm{Q}}$, $\tilde{\mathrm{C}}$ are not shown. - In the column Dγ, i.e. $\mathrm{D}\gamma(\mathbf{r}_1|\mathbf{r}'_1, \mathbf{r}_2|\mathbf{r}'_2) = \gamma(\mathbf{r}_1|\mathbf{r}_1, \mathbf{r}_2|\mathbf{r}_2)$ the PDs $g_{\uparrow\uparrow}$, $g_{\downarrow\downarrow}$, g_a, and Δg are defined. They depend on $r_{12} = |\mathbf{r}_1 - \mathbf{r}_2|$, whereas all the position matrices depend on $\mathbf{r}_1|\mathbf{r}'_1, \mathbf{r}_2|\mathbf{r}'_2$. Note $\mathrm{D}\gamma_a^1 = \mathrm{D}\gamma_a^2$, which holds because of $\langle \hat\psi_\uparrow^\dagger(\mathbf{r}_1)\hat\psi_\downarrow^\dagger(\mathbf{r}_2)\hat\psi_\downarrow(\mathbf{r}_2)\hat\psi_\uparrow(\mathbf{r}_1)\rangle = \langle \hat\psi_\downarrow^\dagger(\mathbf{r}_2)\hat\psi_\uparrow^\dagger(\mathbf{r}_1)\hat\psi_\uparrow(\mathbf{r}_1)\hat\psi_\downarrow(\mathbf{r}_2)\rangle$ and $r_{21} = r_{12}$. It is similarly $\mathrm{D}\gamma_a^3 = \mathrm{D}\gamma_a^4$, but for the off-diagonal elements it is $\gamma_a^1 \neq \gamma_a^2$ and $\gamma_a^3 \neq \gamma_a^4$. Note further, that the spin-antiparallel 2-matrices γ_a^1, γ_a^2, and γ_a^3, γ_a^4 are indeed hermitian, but have not the Pauli-principle property $\mathrm{P}\gamma = \pm\gamma$.

To calculate V_{ss} and V_a we need only the diagonal elements $\mathrm{D}\gamma_{ss}$ and $D\gamma_a^{1,2}$. Also for the contractions only $\mathrm{C}\gamma_{ss}$ and $\mathrm{C}\gamma_a^{1,2}$ are needed. So the question arises: what is the physical meaning of the spin-exchange matrices $\gamma_a^{3,4}$, which information they contain ? Because of $\tilde{\mathrm{D}}\gamma_a^3 = -\mathrm{D}\gamma_a^1$, $\tilde{\mathrm{D}}\gamma_a^4 = -\mathrm{D}\gamma_a^2$ they contain in principle the same PD-information as $\gamma_a^{1,2}$. The 2-matrices $\gamma_a^{3,4}$ mimick spin-flip processes such that the total spin is conserved. Thus they play a role only if the particle interaction would be spin-dependent, e.g. different for singlet and triplet spin-states. The normalization of $\gamma_a^{3,4}$, $\mathrm{Tr}\gamma_a^{3,4} = -Na(r_s, \zeta)$, does not appear in the normalization of the total 2-matrix γ_2, because of $\mathrm{Tr}\delta_a^{3,4} = 0$ as shown below in Eq.(22). For the special cases 'no interaction' and 'no polarization' cf. Apps. 1 and 2.

With the spin-matrices (corresponding to the six matrix elements $\gamma_{ss}, \gamma_a^1, \cdots, \gamma_a^4$, namely)

$$\begin{aligned}
\delta_{\uparrow\uparrow}(\sigma_1|\sigma'_1, \sigma_2|\sigma'_2) &= \delta_{\sigma_1,\sigma'_1}\ \delta_{\sigma_2,\sigma'_2}\delta_{\sigma_1,\uparrow}\delta_{\sigma_2,\uparrow}, && \mathrm{Tr}\delta_{\uparrow\uparrow} = 1, \\
\delta_{\downarrow\downarrow}(\sigma_1|\sigma'_1, \sigma_2|\sigma'_2) &= \delta_{\sigma_1,\sigma'_1}\ \delta_{\sigma_2,\sigma'_2}\delta_{\sigma_1,\downarrow}\delta_{\sigma_2,\downarrow}, && \mathrm{Tr}\delta_{\downarrow\downarrow} = 1, \\
\delta_a^1(\sigma_1|\sigma'_1, \sigma_2|\sigma'_2) &= \delta_{\sigma_1,\sigma'_1}\ \delta_{\sigma_2,\sigma'_2}\delta_{\sigma_1,\uparrow}\delta_{\sigma_2,\downarrow}, && \mathrm{Tr}\delta_a^1 = 1, \\
\delta_a^2(\sigma_1|\sigma'_1, \sigma_2|\sigma'_2) &= \delta_{\sigma_1,\sigma'_1}\ \delta_{\sigma_2,\sigma'_2}\delta_{\sigma_1,\downarrow}\delta_{\sigma_2,\uparrow}, && \mathrm{Tr}\delta_a^2 = 1, \\
\delta_a^3(\sigma_1|\sigma'_1, \sigma_2|\sigma'_2) &= \delta_{\sigma_1,\sigma'_2}\ \delta_{\sigma_2,\sigma'_1}\delta_{\sigma_1,\uparrow}\delta_{\sigma_2,\downarrow}, && \mathrm{Tr}\delta_a^3 = 0, \\
\delta_a^4(\sigma_1|\sigma'_1, \sigma_2|\sigma'_2) &= \delta_{\sigma_1,\sigma'_2}\ \delta_{\sigma_2,\sigma'_1}\delta_{\sigma_1,\downarrow}\delta_{\sigma_2,\uparrow}, && \mathrm{Tr}\delta_a^4 = 0
\end{aligned} \tag{22}$$

(note $\mathrm{D}\delta_a^{3,4} = 0$) the total 2-matrix γ_2 is given by

$$\begin{aligned}
\gamma_2 &= \delta_{\uparrow\uparrow}\gamma_{\uparrow\uparrow} + \delta_{\downarrow\downarrow}\gamma_{\downarrow\downarrow} + \delta_a^1\gamma_a^1 + \delta_a^2\gamma_a^2 + \delta_a^3\gamma_a^3 + \delta_a^4\gamma_a^4, \\
\mathrm{Tr}\gamma_2 &= N_\uparrow(N_\uparrow - 1) + N_\downarrow(N_\downarrow - 1) + N_\uparrow N_\downarrow + N_\downarrow N_\uparrow + 0 + 0 = N(N-1). \qquad (23)
\end{aligned}$$

From Eq.(23) also follows the spin-traced 2-matrix $\gamma_{\uparrow\uparrow} + \gamma_{\downarrow\downarrow} + 2\gamma_a$, where $\gamma_a = \frac{1}{2}(\gamma_a^1 + \gamma_a^2)$.

In view of the spectral resolution of γ_2 (respectively of its position matrices) in terms of antisymmetric spin geminals, the drawback of Eq.(23) is that $\gamma_a^1, \cdots, \gamma_a^4$ are not eigenfunctions of the permutation operator P as seen from Table I, column $\mathrm{P}\gamma$. Only the linear combinations $\gamma_\pm$ have the eigenvalues $\pm$. So it is useful to rewrite Eq.(23) by appropriate (the Pauli symmetry achieving) linear combinations. In the first (intermediate) step with the definitions of Table I for δ_a, γ_a etc. (the last but one block of lines) it results the parallel/antiparallel representation

$$\begin{aligned}
\gamma_2 &= \delta_{\uparrow\uparrow}\gamma_{\uparrow\uparrow} + \delta_{\downarrow\downarrow}\gamma_{\downarrow\downarrow} + \delta_a\gamma_a + \delta_a'\gamma_a' + \tilde{\delta}_a\tilde{\gamma}_a + \tilde{\delta}_a'\tilde{\gamma}_a', \\
\mathrm{Tr}\gamma_2 &= N_\uparrow(N_\uparrow - 1) + N_\downarrow(N_\downarrow - 1) + 2N_\uparrow N_\downarrow + 0 + 0 + 0 = N(N-1). \qquad (24)
\end{aligned}$$

In the next step using $\delta_\pm, \gamma_\pm$ etc. of Table I (the last block of lines) it turns out

$$\begin{aligned}
\gamma_2 &= [\delta_{\uparrow\uparrow}\gamma_{\uparrow\uparrow} + \delta_{\downarrow\downarrow}\gamma_{\downarrow\downarrow} + \delta_+\gamma_-] + \delta_-\gamma_+ + [\tilde{\delta}_-\tilde{\gamma}_+ + \tilde{\delta}_+\tilde{\gamma}_-], \\
\mathrm{Tr}\gamma_2 &= [N_\uparrow(N_\uparrow - 1) + N_\downarrow(N_\downarrow - 1) + (N_\uparrow N_\downarrow - Na)] + (N_\downarrow N_\uparrow + Na) + [0+0] = N(N-1).
\end{aligned}$$
$$(25)$$

This is the triplet/singlet representation of the 2-matrix γ_2, because the first three terms describe the spin-triplet, the next term the spin-singlet, and the last two terms a singlet/triplet mixing, cf. App. 4. Note, that in the trace the terms $\pm Na$ and in the diagonal elements the terms $\pm\rho_\uparrow\rho_\downarrow\Delta g$ cancel each other. The symmetrized, respectively antisymmetrized spin-matrices used in Eq.(25) are

$$\begin{aligned}
\delta_\pm(\sigma_1|\sigma_1', \sigma_2|\sigma_2') &= \frac{1}{2}(\delta_{\sigma_1,\sigma_1'}\ \delta_{\sigma_2,\sigma_2'} \pm \delta_{\sigma_1,\sigma_2'}\ \delta_{\sigma_2,\sigma_1'})(\delta_{\sigma_1\uparrow}\delta_{\sigma_2\downarrow} + \delta_{\sigma_1\downarrow}\delta_{\sigma_2\uparrow}), \quad \mathrm{Tr}\delta_\pm = 1, \\
\tilde{\delta}_\pm(\sigma_1|\sigma_1', \sigma_2|\sigma_2') &= \frac{1}{2}(\delta_{\sigma_1,\sigma_1'}\ \delta_{\sigma_2,\sigma_2'} \mp \delta_{\sigma_1,\sigma_2'}\ \delta_{\sigma_2,\sigma_1'})(\delta_{\sigma_1\uparrow}\delta_{\sigma_2\downarrow} - \delta_{\sigma_1\downarrow}\delta_{\sigma_2\uparrow}), \quad \mathrm{Tr}\tilde{\delta}_\pm = 0 \qquad (26)
\end{aligned}$$

and the corresponding position matrices are

$$\begin{aligned}
\gamma_\pm &= \frac{1}{2}(\gamma_a^1 + \gamma_a^2 \mp \gamma_a^3 \mp \gamma_a^4), \quad \mathrm{Tr}\gamma_\pm = N_\uparrow N_\downarrow \pm Na, \\
\tilde{\gamma}_\pm &= \frac{1}{2}(\gamma_a^1 - \gamma_a^2 \pm \gamma_a^3 \mp \gamma_a^4), \quad \mathrm{Tr}\tilde{\gamma}_\pm = 0, \qquad (27)
\end{aligned}$$

as it follows from Table I.

Hermiticity: Whereas $\gamma_{\uparrow\uparrow}$, $\gamma_{\downarrow\downarrow}$, and $\gamma_\pm$ are hermitian, the last two terms of Eq.(25) are hermitian only summed together because of $Q\tilde{\delta}_\pm = \tilde{\delta}_\mp$ and $Q\tilde{\gamma}_\pm = \tilde{\gamma}_\mp$. - As above the spin-traced 2-matrix is again $\gamma_{\uparrow\uparrow} + \gamma_{\downarrow\downarrow} + 2\gamma_a$, because of $\gamma_+ + \gamma_- = \gamma_a^1 + \gamma_a^2 = 2\gamma_a$. - Note that the interaction energy V consists of a triplet contribution $V_{\uparrow\uparrow} + V_{\downarrow\downarrow} + V_-$ arising from γ_{ss}, γ_- and of a singlet contribution V_+ stemming from γ_+.

6. The pair densities

For the diagonal elements $\gamma_2(1|1,2|2) = \rho_2(1,2) = \rho^2 g(1,2)$, from Eq.(24) and Eq.(25) the two equivalent expressions ($r = r_{12}$)

$$\begin{aligned}\rho^2 g(1,2) &= \delta_{\sigma_1\uparrow}\delta_{\sigma_2\uparrow}\rho_\uparrow^2 g_{\uparrow\uparrow}(r) + \delta_{\sigma_1\downarrow}\delta_{\sigma_2\downarrow}\rho_\downarrow^2 g_{\downarrow\downarrow}(r) + \delta_{\sigma_1,-\sigma_2}\rho_\uparrow\rho_\downarrow g_a(r) \quad \text{or} \\ \rho^2 g(1,2) &= \left[\delta_{\sigma_1\uparrow}\delta_{\sigma_2\uparrow}\rho_\uparrow^2 g_{\uparrow\uparrow}(r) + \delta_{\sigma_1\downarrow}\delta_{\sigma_2\downarrow}\rho_\downarrow^2 g_{\downarrow\downarrow}(r) + \frac{1}{2}\delta_{\sigma_1,-\sigma_2}\rho_\uparrow\rho_\downarrow g_-(r)\right] + \frac{1}{2}\delta_{\sigma_1,-\sigma_2}\rho_\uparrow\rho_\downarrow g_+(r)\end{aligned} \tag{28}$$

result. The two dimensionless spin-parallel PDs $g_{\uparrow\uparrow}(r), g_{\downarrow\downarrow}(r)$ are (with $\mathbf{r}_1' = \mathbf{r}_1$, $\mathbf{r}_2' = \mathbf{r}_2$) defined and with $\mathrm{Tr}\gamma_{ss} = N_s(N_s - 1)$ normalized by

$$\begin{aligned}\rho_\uparrow^2 g_{\uparrow\uparrow}(r) &= \gamma_{\uparrow\uparrow}(\mathbf{r}_1|\mathbf{r}_1, \mathbf{r}_2|\mathbf{r}_2), \ \int d^3r\ \rho[1 - g_{\uparrow\uparrow}(r)] = \frac{2}{1+\zeta}, \ g_{\uparrow\uparrow}(\infty) = 1, \\ \rho_\downarrow^2 g_{\downarrow\downarrow}(r) &= \gamma_{\downarrow\downarrow}(\mathbf{r}_1|\mathbf{r}_1, \mathbf{r}_2|\mathbf{r}_2), \ \int d^3r\ \rho[1 - g_{\downarrow\downarrow}(r)] = \frac{2}{1-\zeta}, \ g_{\downarrow\downarrow}(\infty) = 1.\end{aligned} \tag{29}$$

The spin-antiparallel PDs $g_a(r)$, $g_\pm(r)$ are defined and with $\mathrm{Tr}\gamma_a = N_\uparrow N_\downarrow$, $\mathrm{Tr}\gamma_\pm = N_\uparrow N_\downarrow \pm Na$ normalized as

$$\begin{aligned}\rho_\uparrow\rho_\downarrow g_a(r) &= \gamma_a(\mathbf{r}_1|\mathbf{r}_1, \mathbf{r}_2|\mathbf{r}_2), \ \int d^3r\ \rho[1 - g_a(r)] = 0, \quad g_a(\infty) = 1, \\ \rho_\uparrow\rho_\downarrow g_\pm(r) &= \gamma_\pm(\mathbf{r}_1|\mathbf{r}_1, \mathbf{r}_2|\mathbf{r}_2), \ \int d^3r\ \rho[1 - g_\pm(r)] = \mp\frac{4a}{1-\zeta^2}, \quad g_\pm(\infty) = 1,\end{aligned} \tag{30}$$

where $4a/(1-\zeta^2)$ equals $2/(1+\zeta)$ for 'no interaction' and 2 for 'no polarization', cf. Apps. 1 and 2. g_+ and g_- are the singlet and triplet spin-antiparallel PDs, respectively. (Here they are normalized as $g_\pm(\infty) = 1$, whereas in Refs. [21, 22] it is $g_\pm(\infty) = 1/2$.) For spin-independent particle interaction the PDs g_+ and g_- are equally weighted, so only their sum $g_+ + g_- = 2g_a$ appears (and not $\frac{1}{2}(g_+ - g_-) = \Delta g$, which is the singlet-triplet splitting of the PDs g_+ and g_-). Spin summation yields the spin-weighted PD $g(r) = \sum_{\sigma_{1,2}} g(1,2)$, which is also given only by g_{ss} and g_a:

$$\begin{aligned} g(r) &= (\frac{\rho_\uparrow}{\rho})^2 g_{\uparrow\uparrow}(r) + (\frac{\rho_\downarrow}{\rho})^2 g_{\downarrow\downarrow}(r) + 2\frac{\rho_\uparrow\rho_\downarrow}{\rho^2} g_a(r) \quad \text{or} \\ 1 - g(r) &= (\frac{\rho_\uparrow}{\rho})^2 [1 - g_{\uparrow\uparrow}(r)] + (\frac{\rho_\downarrow}{\rho})^2 [1 - g_{\downarrow\downarrow}(r)] + 2\frac{\rho_\uparrow\rho_\downarrow}{\rho^2}[1 - g_a(r)] \end{aligned} \tag{31}$$

with $\rho_\uparrow/\rho = (1+\zeta)/2$ and $\rho_\downarrow/\rho = (1-\zeta)/2$ and $\int d^3r\ \rho[1 - g(r)] = 1, g(\infty) = 1$.

'On-top', i.e. for $r = 0$, the coalescing cusp theorems

$$g'''_{ss}(0) = \frac{3}{2}\alpha_0 g''_{ss}(0), \quad g'_a(0) = \alpha_0 g_a(0), \quad \alpha_0 = \left(\frac{4}{9\pi}\right)^{1/3} r_s \tag{32}$$

hold [17, 33].

7. The contraction SRs

The contraction of the spin-parallel 2-matrices,

$$\begin{aligned} &\int d^3r_2 \gamma_{ss}(\mathbf{r}_1|\mathbf{r}'_1, \mathbf{r}_2|\mathbf{r}_2) = \gamma_s(\mathbf{r}_1|\mathbf{r}'_1)(N_s - 1) \quad \text{or} \\ &\int d^3r_2 [\rho_s \gamma_s(\mathbf{r}_1|\mathbf{r}'_1) - \gamma_{ss}(\mathbf{r}_1|\mathbf{r}'_1, \mathbf{r}_2|\mathbf{r}_2)] = \gamma_s(\mathbf{r}_1|\mathbf{r}'_1), \end{aligned} \tag{33}$$

makes the spin-parallel 3-point functions $\gamma_{ss}(\mathbf{r}_1|\mathbf{r}'_1, \mathbf{r}_2|\mathbf{r}_2)$ to appear. Because of translational and rotational invariance (six conditions) this function depends only on the three distances $r_{11'} = |\mathbf{r}_1 - \mathbf{r}'_1|$, $r_{12} = |\mathbf{r}_1 - \mathbf{r}_2|$, and $r_{1'2} = |\mathbf{r}'_1 - \mathbf{r}_2|$, besides the direction of $\mathbf{r}_{12}$ can be averaged:

$$G_{ss}(r_{11'}, r_{12}) = \int \frac{d\Omega_{12}}{4\pi} G_{ss}(r_{11'}, r_{12}, |\mathbf{r}_{11'} - \mathbf{r}_{12}|), \quad \rho_s^2 G_{ss}(r_{11'}, r_{12}, r_{1'2}) = \gamma_{ss}(\mathbf{r}_1|\mathbf{r}'_1, \mathbf{r}_2|\mathbf{r}_2). \tag{34}$$

$G_{\uparrow\uparrow}(r_{11'}, r_{12})$ and $G_{\downarrow\downarrow}(r_{11'}, r_{12})$ are special 2-matrices yielding with $r'_1 = r_1$ the spin-parallel PDs: $g_{ss}(r_{12}) = G_{ss}(0, r_{12})$. The contraction SRs (33) in terms of these 2-matrices $G_{ss}(r_{11'}, r_{12})$ are rewritten as

$$\int d^3r_{12}\ \rho_s[f_s(r_{11'}) - G_{ss}(r_{11'}, r_{12})] = f_s(r_{11'}), \quad G_{ss}(r_{11'}, \infty) = f_s(r_{11'}). \tag{35}$$

So the functions G_{ss} contain not only $g_{ss}(r)$, but also $f_s(r)$ and with $r_{11'} = 0$ the SRs (35) yield the normalization (29) of $g_{ss}(r)$.

The contraction of the spin-antiparallel 2-matrices $\gamma_\pm$ yields

$$\int d^3r_2 \gamma_\pm(\mathbf{r}_1|\mathbf{r}_1', \mathbf{r}_2|\mathbf{r}_2) = \frac{1}{2}[N_\downarrow \gamma_\uparrow(\mathbf{r}_1|\mathbf{r}_1') + N_\uparrow \gamma_\downarrow(\mathbf{r}_1|\mathbf{r}_1')] \pm \rho\alpha(r_{11'}) \quad \text{or,}$$

$$\int d^3r_2 \left\{ \frac{1}{2}[\rho_\downarrow \gamma_\uparrow + \rho_\uparrow \gamma_\downarrow](\mathbf{r}_1|\mathbf{r}_1') - \gamma_a(\mathbf{r}_1|\mathbf{r}_1', \mathbf{r}_2|\mathbf{r}_2) \right\} = \mp \rho\alpha(r_{11'}). \tag{36}$$

Corresponding 2-point functions are defined by

$$\rho_\uparrow \rho_\downarrow G_\pm(r_{11'}, r_{12}) = \int \frac{d\Omega_{12}}{4\pi} \gamma_\pm(\mathbf{r}_1|\mathbf{r}_1', \mathbf{r}_2|\mathbf{r}_2). \tag{37}$$

With this definition of $G_\pm$ the contraction SRs (36) can be rewritten as

$$\int d^3r_{12}\ \rho[f(r_{11'}) - G_\pm(r_{11'}, r_{12})] = \mp \frac{4\alpha(r_{11'})}{1-\zeta^2}, \quad G_\pm(r_{11'}, \infty) = f(r_{11'}) \tag{38}$$

with $f(r) = \frac{1}{2}[f_\uparrow(r) + f_\downarrow(r)]$. The functions $G_\pm$ contain the spin-antiparallel PDs $g_\pm(r_{12}) = G_\pm(0, r_{12})$. Thus the SRs (38) contain with $r_{11'} = 0$ the normalizations (30) of $g_\pm(r)$. Note that the 1-matrix $\alpha(r_{11'})$ determines the normalization of the PDs $g_\pm$ with $\alpha(0) = a$. - For $G_a = (G_+ + G_-)/2$, which equally weights the singlet/triplet contributions, it holds

$$\int d^3r_{12}\ \rho[f(r_{11'}) - G_a(r_{11'}, r_{12})] = 0, \quad G_a(r_{11'}, \infty) = f(r_{11'}). \tag{39}$$

In summary, the two functions $G_{ss}(r_{11'}, r_{12})$ not only contain the two spin-parallel PDs $g_{ss}(r) = G_{ss}(0, r)$, but also the two 1-matrices $f_s(r) = G_{ss}(r, \infty)$. And the function $G_a(r_{11'}, r_{12})$ contains the spin-antiparallel PD $g_a(r) = G_a(0, r)$ and besides it controls $f = \frac{1}{2}[f_\uparrow + f_\downarrow]$ with $\frac{1}{2}[f_\uparrow(r) + f_\downarrow(r)] = G_a(r, \infty)$. A more refined description distinguishes between the singlet function G_+ and the triplet function G_-. So there are six SRs, namely the four contraction SRs (35) and (38) and the two momentum-distribution normalizations $f_s(0) = 1$.

8. Wick's theorem, linked diagrams, and the cumulant expansion

If the Coulomb repulsion is treated as perturbation (with many-body S-matrix theory and Wick's theorem), then the 1-matrices $\gamma_\uparrow$ and $\gamma_\downarrow$ are given by *linked* Feynman diagrams only, as sketched in (Eq.(15) shows with $\gamma_1 \sim \delta_{\sigma_1,\sigma_2}$: there is no spin-flip along an electron line)

$\langle \hat{\psi}_s^\dagger(\mathbf{r}_1')\hat{\psi}_s(\mathbf{r}_1)\rangle =$  $+$  $+ \cdots .$ (40)

Contrary to this the 2-matrices are given by *both* unlinked (or disconnected) *and* linked (or connected) diagrams. If all the unlinked diagrams are summed up, products of the 1-matrices appear, e.g.

$$\gamma_{ss} = \langle ssss\rangle = \gamma_s(\mathbf{r}_1|\mathbf{r}_1')\gamma_s(\mathbf{r}_2|\mathbf{r}_2') - \gamma_s(\mathbf{r}_1|\mathbf{r}_2')\gamma_s(\mathbf{r}_2|\mathbf{r}_1') + \langle ssss\rangle_{\mathrm{c}}, \tag{41}$$

where $\langle ssss\rangle_c = -\chi_{ss}$ is given by the sum of all the connected diagrams of

$$\chi_{ss} = \langle \hat{\psi}_s^\dagger(\mathbf{r}_1')\hat{\psi}_s^\dagger(\mathbf{r}_2')\hat{\psi}_s(\mathbf{r}_2)\hat{\psi}_s(\mathbf{r}_1)\rangle_c =$$



$$+$$



$$+\cdots . \tag{42}$$

The index ‘c’ means ‘connected’ or ‘cumulant’. It is similarly

$$\begin{aligned}\gamma_a^1 &= \langle\uparrow\downarrow\downarrow\uparrow\rangle = \gamma_\uparrow(\mathbf{r}_1|\mathbf{r}_1')\gamma_\downarrow(\mathbf{r}_2|\mathbf{r}_2') + \langle\uparrow\downarrow\downarrow\uparrow\rangle_{\mathrm{c}},\\ \gamma_a^3 &= \langle\downarrow\uparrow\downarrow\uparrow\rangle = \gamma_\uparrow(\mathbf{r}_1|\mathbf{r}_2')\gamma_\downarrow(\mathbf{r}_2|\mathbf{r}_1') + \langle\downarrow\uparrow\downarrow\uparrow\rangle_{\mathrm{c}},\end{aligned} \tag{43}$$

where $\langle\uparrow\downarrow\downarrow\uparrow\rangle_{\mathrm{c}} = -\chi_a^1$ and $\langle\downarrow\uparrow\downarrow\uparrow\rangle_{\mathrm{c}} = -\chi_a^3$ mean the linked diagrams of

$$\chi_a^1 = \langle \hat{\psi}_\uparrow^\dagger(\mathbf{r}_1')\hat{\psi}_\downarrow^\dagger(\mathbf{r}_2')\hat{\psi}_\downarrow(\mathbf{r}_2)\hat{\psi}_\uparrow(\mathbf{r}_1)\rangle_c =$$



$$+\ \cdots ,$$

$$\chi_a^3 = \langle \hat{\psi}_\downarrow^\dagger(\mathbf{r}_1')\hat{\psi}_\uparrow^\dagger(\mathbf{r}_2')\hat{\psi}_\downarrow(\mathbf{r}_2)\hat{\psi}_\uparrow(\mathbf{r}_1)\rangle_c =$$



$$+\ \cdots . \tag{44}$$

The products of the 1-matrices $\gamma_s = \rho_s f_s$ may be called generalized Hartree-Fock (HF) parts, where the term ‘generalized’ means ‘built up from the true, hence non-idempotent 1-matrix $f_s(r)$’. The HF parts of the six position matrices (21) together with eight other related 2-matrices and their properties are listed in Table II. The matrices χ_{ss}, χ_a^1, χ_a^3 etc. are called cumulant 2-matrices, they are listed in Table III. Eqs.(41) and (43) are the first steps of the more general cumulant expansion [34]. Per construction, the cumulant matrices are given by linked diagrams, what makes them size-extensively normalizable. Contrary to this the normalizations of the γ-matrices and the $\gamma^{\rm HF}$-matrices contain terms $\sim N^2$, cf. the columns 'Tr' of Table III with those of Tables I and II. In the column C$\gamma^{\rm HF}$ of Table II the already above introduced 1-matrices $\beta_s(r_{11'})$ and $\beta_a(r_{11'})$ appear, as it is obvious from Eq.(19). From the diagonal elements D$\gamma^{\rm HF}$ follow the PDs $g_{ss}^{\rm HF}(r) = 1 - |f_s(r)|^2$ with $\rho_s \int d^3r[1 - g_{ss}^{\rm HF}(r)] = \frac{N}{N_s} b_s$ and $g_a^{\rm HF}(r) = 1$ as well as $\Delta g^{\rm HF}(r) = f_\uparrow(r) f_\downarrow(r)$.

The diagonal elements of χ_{ss} and χ_a^1 or χ_a^2 give the cumulant contributions to the PDs:

$$\begin{aligned}
\rho_s^2 h_{ss} &= {\rm D}\chi_s, \int d^3r\ \rho h_{ss}(r) = \frac{\rho}{\rho_s}(1 - \frac{\rho}{\rho_s} b_s),\ g_{ss}(r) = 1 - |f_s(r)|^2 - h_s(r), \\
\rho_\uparrow \rho_\downarrow h_\pm &= {\rm D}\chi_\pm, \int d^3r\ \rho h_\pm(r) = \mp(a - b_a), g_\pm(r) = 1 \mp [\alpha(r) - \beta_a(r)] - h_\pm(r), \\
\rho_\uparrow \rho_\downarrow h_a &= {\rm D}\chi_a, \int d^3r\ \rho h_a(r) = 0,\ g_a(r) = 1 - 0 - h_a(r).
\end{aligned} \tag{45}$$

The contraction of the cumulant 2-matrix: With the definitions

$$\begin{aligned}
\rho_s^2 H_{ss}(r_{11'}, r_{12}) &= \int \frac{d\Omega_{12}}{4\pi} \int d^3r_2 \chi_{ss}(\mathbf{r}_1|\mathbf{r}_1', \mathbf{r}_2|\mathbf{r}_2), \\
\rho_\uparrow \rho_\downarrow H_\pm(r_{11'}, r_{12}) &= \int \frac{d\Omega_{12}}{4\pi} \int d^3r_2 \chi_\pm(\mathbf{r}_1|\mathbf{r}_1', \mathbf{r}_2|\mathbf{r}_2),
\end{aligned} \tag{46}$$

and $H_a = (H_+ + H_-)/2$ the contraction SRs are

$$\begin{aligned}
\int d^3r_{12}\ \rho H_{ss}(r_{11'}, r_{12}) &= \frac{\rho}{\rho_s}[f_s(r_{11'}) - \frac{\rho}{\rho_s}\beta_s(r_{11'})], \\
\int d^3r_{12}\ \rho H_\pm(r_{11'}, r_{12}) &= \mp \frac{\rho^2}{\rho_\uparrow \rho_\downarrow}[\alpha(r_{11'}) - \beta_a(r_{11'})], \\
\int d^3r_{12}\ \rho H_a(r_{11'}, r_{12}) &= 0.
\end{aligned} \tag{47}$$

The advantage of the H-functions: in perturbation theory they are given by linked diagrams, which are size-extensive. They contain the cumulant PDs $h_{ss}(r) = H_{ss}(0, r)$, $h_\pm(r) = H_\pm(0, r)$, $h_a(r) = H_a(0, r)$. To get also the 1-matrices f_s (or equivalently the momentum distributions n_s) one has to write the first equation of the SRs (46) as

$$\int d^3r_{12}\ \rho H_{ss}(r_{11'}, r_{12}) = \left(\frac{\rho}{\rho_s}\right)^2 \frac{1}{N} \sum_{\mathbf{k}} n_s(k)[1 - n_s(k)] {\rm e}^{i\mathbf{k}\mathbf{r}_{11'}} \tag{48}$$

and to Fourier analyse the lhs according to

$$\int d^3r_{12}\ \rho H_{ss}(r_{11'}, r_{12}) = \left(\frac{\rho}{\rho_s}\right)^2 \frac{1}{N} \sum_{\mathbf{k}} m_s(k) {\rm e}^{i\mathbf{k}\mathbf{r}_{11'}} \tag{49}$$

defining functions $m_s(k)$. Then one has to solve the quadratic equation $n_s^2(k) - n_s(k) + m_s(k) = 0$. As already mentioned, one way to get the PDs and the momentum distributions is to perturbatively calculate linked diagrams for H_{ss} and $H_\pm$. Another way is to solve an effective 2-body Schrödinger equation for the geminals, which parametrize the 2-matrices γ_{ss} and $\gamma_\pm$ like this has been done successfully for the spin-unpolarized HEG [17, 18, 19, 20]. This way is sketched in the following for the spin-polarized HEG.

9. The spectral resolutions

The four 2-matrices γ_{ss} and $\gamma_\pm$ are needed for the key functions G_{ss} and $G_\pm$. They have the correct Pauli symmetry and are hermitian. So they can be diagonalized in terms of symmetric/antisymmetric geminals and corresponding occupancies. This is similar to the well-known natural representation of the 1-matrix. For the HF parts this diagonalization can be done easily with the help of the spectral resolution or natural-orbital representation (17) of the 1-matrices $f_s(r)$ in terms of plane waves (because the system is homogeneous). Thereby the geminals

$$\frac{1}{\sqrt{\Omega}}\mathrm{e}^{\mathrm{i}\mathbf{K}\mathbf{R}}\frac{1}{\sqrt{\Omega}}\mathrm{e}^{\mathrm{i}\mathbf{k}\mathbf{r}} \quad \text{with} \quad \mathbf{R} = \frac{1}{2}(\mathbf{r}_1 + \mathbf{r}_2),\ \mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2 \tag{50}$$

appear together with the weights

$$\begin{gathered}\mu_s(\mathbf{K},\mathbf{k}) = n_s(k_1)n_s(k_2)\ , \qquad \mu_a(\mathbf{K},\mathbf{k}) = n_\uparrow(k_1)n_\downarrow(k_2)\\ \text{with} \quad \mathbf{k}_1 = \frac{1}{2}\mathbf{K} + \mathbf{k}\ ,\ \ \mathbf{k}_2 = \frac{1}{2}\mathbf{K} - \mathbf{k}.\end{gathered} \tag{51}$$

$\mathbf{R}$ is the centre-of-mass coordinate and $\frac{1}{\sqrt{\Omega}}\mathrm{e}^{\mathrm{i}\mathbf{K}\mathbf{R}}$ describes the centre-of-mass motion (which is a free-particle motion, because the system is homogeneous), whereas $\mathbf{r}$ is the relative coordinate and $\varphi^0(\mathbf{r},\mathbf{k}) = \frac{1}{\sqrt{\Omega}}\mathrm{e}^{\mathrm{i}\mathbf{k}\mathbf{r}}$ describes the relative motion of the HF parts. Starting with the expressions for $\gamma_{ss}^{\mathrm{HF}}$ and $\gamma_\pm^{\mathrm{HF}}$ of Table II, the following spectral resolutions of the HF parts result

$$\begin{aligned}\gamma_{ss}^{\mathrm{HF}}(\mathbf{R}|\mathbf{R}',\mathbf{r}|\mathbf{r}') &= \sum_{\mathbf{K},\mathbf{k}} \mu_s(\mathbf{K},\mathbf{k})\varphi_-^0(\mathbf{R},\mathbf{r})\varphi_-^{0*}(\mathbf{R}',\mathbf{r}'),\\ \gamma_\pm^{\mathrm{HF}}(\mathbf{R}|\mathbf{R}',\mathbf{r}|\mathbf{r}') &= \sum_{\mathbf{K},\mathbf{k}} \mu_a(\mathbf{K},\mathbf{k})\varphi_\pm^0(\mathbf{R},\mathbf{r})\varphi_\pm^{0*}(\mathbf{R}',\mathbf{r}')\end{aligned} \tag{52}$$

with $\varphi_\pm^0(\mathbf{R},\mathbf{r}) = \frac{1}{\sqrt{\Omega}}\mathrm{e}^{\mathrm{i}\mathbf{K}\mathbf{R}}\frac{4\pi}{\sqrt{\Omega}}\sum_L^\pm \mathrm{i}^l j_L(k\mathbf{r})Y_L^*(\mathbf{e}_k)$, where the dependence of $\mathbf{K},\mathbf{k}$ is not explicitly denoted and the angular momentum expansion of a plane wave is used. Note $L = (l, m_l)$

and $j_L(k\mathbf{r}) = j_l(kr)Y_L(\mathbf{e}_r)$. So the spectral resolution (52) can be written in terms of free-electron geminals $j_l(kr)$ and occupancy matrices $\mu^s_{LL'}(\mathbf{K}, k)$ and $\mu^a_{LL'}(\mathbf{K}, k)$. The normalizations

$$\begin{aligned}
\mathrm{Tr}\gamma_{ss}^{\mathrm{HF}} &= \sum_{\mathbf{k}_{1,2}} \mu_s(\mathbf{K},\mathbf{k})[1-\delta_{\mathbf{k}_1,\mathbf{k}_2}] = N_s^2 - \sum_{\mathbf{k}} n_s^2(k) = N_s^2 - Nb_s,\\
\mathrm{Tr}\gamma_{\pm}^{\mathrm{HF}} &= \sum_{\mathbf{k}_{1,2}} \mu_a(\mathbf{K},\mathbf{k})[1\pm\delta_{\mathbf{k}_1,\mathbf{k}_2}] = N_\uparrow N_\downarrow \pm \sum_{\mathbf{k}} n_\uparrow(k) n_\downarrow(k) = N_\uparrow N_\downarrow \pm Nb_a
\end{aligned} \tag{53}$$

follow from $\int \frac{d^3r}{\Omega}\mathrm{e}^{\mathrm{i}\mathbf{k}\mathbf{r}} = \delta_{\mathbf{k},0}$ in agreement with Table II.

From the spectral resolution (52) of the HF parts one may derive the spectral resolution of the total 2-matrix by replacing the the free-electron geminals $\varphi^0_\pm$ or $j_l(kr)$ by interacting-electron geminals $\varphi_\pm$ or $R_l(r,k)$:

$$\begin{aligned}
\gamma_{ss}(\mathbf{R}|\mathbf{R}',\mathbf{r}|\mathbf{r}') &= \sum_{\mathbf{K},\mathbf{k}} \mu_s(\mathbf{K},\mathbf{k})\varphi_-(\mathbf{R},\mathbf{r})\varphi_-^*(\mathbf{R}',\mathbf{r}'),\\
\gamma_\pm(\mathbf{R}|\mathbf{R}',\mathbf{r}|\mathbf{r}') &= \sum_{\mathbf{K},\mathbf{k}} \mu_a(\mathbf{K},\mathbf{k})\varphi_\pm(\mathbf{R},\mathbf{r})\varphi_\pm^*(\mathbf{R}',\mathbf{r}').
\end{aligned} \tag{54}$$

(The general structure (54) in terms of eigenfunctions and eigenvalues, which diagonalize γ_{ss} and $\gamma_\pm$, is correct, but the use of Eq.(51) for the geminal weights, which is right in Eq.(52), may be here in Eq.(54) only an approximation.) With the angular-momentum expansion of $\varphi_\pm(\mathbf{R},\mathbf{r})$ the PDs g_{ss} and $g_\pm$ are given in terms of the afore mentioned geminals $R_l(r,k)$ and occupancies $\mu^s_{LL'}(\mathbf{K},k)$ and $\mu^a_{LL'}(\mathbf{K},k)$. Their diagonalization causes a $(\mathbf{K},k)$-dependent L-mixing with L-mixed geminals $R^\pm_\Lambda(\mathbf{r},k)$ and occupancies $\mu^\pm_\Lambda$. Calculating the PDs with the non-spherically symmetric functions $|R^\pm_\Lambda(\mathbf{r},k)|^2$, their non-sphericity is averaged out by summing over Λ with $\mu^\pm_\Lambda$. The geminals $R_l(r,k)$ are the scattering state solutions of a 2-body Schrödinger equation with an effective repulsion $v_\pm(r) = \frac{e^2}{r} + \cdots$, which may be in general a non-local interaction potential, but a local one possibly can be a good approximation. Whithin a PDFT the ellipsis result from a functional derivative of the kinetic energy as a functional the PD [35].

Eq.(54) means for the cumulant matrices

$$\begin{aligned}
\chi_{ss}(\mathbf{R}|\mathbf{R}',\mathbf{r}|\mathbf{r}') &= \sum_{\mathbf{K},\mathbf{k}} \mu_s(\mathbf{K},\mathbf{k})[\varphi_-^0(\mathbf{R},\mathbf{r})\varphi_-^{0*}(\mathbf{R}',\mathbf{r}') - \varphi_-(\mathbf{R},\mathbf{r})\varphi_-^*(\mathbf{R}',\mathbf{r}')],\\
\chi_\pm(\mathbf{R}|\mathbf{R}',\mathbf{r}|\mathbf{r}') &= \sum_{\mathbf{K},\mathbf{k}} \mu_a(\mathbf{K},\mathbf{k})[\varphi_\pm^0(\mathbf{R},\mathbf{r})\varphi_\pm^{0*}(\mathbf{R}',\mathbf{r}') - \varphi_\pm(\mathbf{R},\mathbf{r})\varphi_\pm^*(\mathbf{R}',\mathbf{r}')].
\end{aligned} \tag{55}$$

Because the cumulant matrices χ_{ss} and $\chi_\pm$ are hermitian, they can be represented alternatively in their 'own' spectral resolution with cumulant weights $\mu^c_{s,a}$ and cumulant geminals R^c_l being the bound state solutions of another 2-body Schrödinger equation with an effective attraction $v^{\rm c}_\pm(r)$, possibly $v^{\rm c}_\pm(r) = -v_\pm(r)$.

10. Scattering phase shifts

The afore mentioned geminals $R_l(r,k)$ are scattering states with an asymptotical behavior

$$R_l(r,k) \to \frac{1}{kr}\sin[kr - l\frac{\pi}{2} + \eta_l(k)] \tag{56}$$

defining scattering phase shifts $\eta_l(k)$ [$v_\pm(r)$ is assumed to decay stronger than $1/r^2$]. For the case 'no polarization' the PD normalizations

$$\int d^3r\ \rho[1 - g_p(r)] = 2, \quad \int d^3r\ \rho[1 - g_a(r)] = 0 \tag{57}$$

can be equivalently reformulated as the Friedel like SR [21]

$$\frac{2}{\pi}\sum\nolimits_L^{\pm}\int_0^\infty dk[-\mu'(k)]\eta_l(k) = \pm c, \quad \mu(k) = \frac{1}{N}\sum_{\mathbf{K}} n(k_1)n(k_2) \tag{58}$$

with $k_{1,2} = |\frac{1}{2}\mathbf{K} \pm \mathbf{k}|$. Generalizing this to the case of non-vanishing polarization, one should expect e.g. for the spin-parallel PDs

$$\frac{2}{\pi}\sum\nolimits_L^{-}\int_0^\infty dk[-\mu_s'(k)]\eta_l(k) = -c_s, \quad \mu_s(k) = \frac{1}{N_s}\sum_{\mathbf{K}} n_s(k_1)n_s(k_2) \tag{59}$$

as an equivalent reformulation of the normalization (29). $3k^2\mu_s(k)$ is the probability of finding two electrons with spin s and momenta $\mathbf{k}_1$ and $\mathbf{k}_2$ with the half momenta difference $k = \frac{1}{2}|\mathbf{k}_1 - \mathbf{k}_2|$ [36]. Similar SRs should hold for electron pairs with antiparallel spins. Also the contraction SRs (29) of Ref. [22] can be generalized to the case of non-vanishing spin polarization.

Summary and outlook

Based on the spin-structure of the 2-matrix [Eq.(25)], its contraction properties (33) and (36) lead to three two-variable functions $G_{\uparrow\uparrow}(x,y)$, $G_{\downarrow\downarrow}(x,y)$, and $G_a(x,y)$, which are key functions, because they contain not only the PDs according to $g_{ss}(r) = G_{ss}(0,r)$ and $g_a(r) = G_a(0,r)$ with $g_{ss}(0) = 0$, $g_a(0) < 1$ and $g_{ss}(r), g_a(r) \geq 0$, but also the 1-matrices according to $f_s(r) = G_{ss}(r, \infty)$ with $f_s(0) = 1$. These key functions G_{ss} and G_a have to obey the contraction SRs (35) and (39). They can be approximately calculated (i) perturbatively, cf. Sec. 8 and Ref. [38] or (ii) by the solution of an effective 2-body equation according to the Kimball-Overhauser approach [18, 19, 20]. The Friedel like normalization and contraction SRs of Refs. [21, 22] can be generalized to the case of non-vanishing spin-polarization. The general analysis presented here has to be completed by numerical studies. Also the question should be studied, whether conclusions can be drawn for the afore mentioned key functions and geminals from the hierarchy of contracted Schrödinger equations [13], the Kummer variety [14], linear inequalities for density matrices [15], the P/D, Q, and G positivity conditions [39, 40]. Future work should concern also the generalization of the fluctuation analysis of the spin-unpolarized HEG [32] to the case of non-vanishing spin-polarization.

Acknowledgments

The authors thank P. Gori-Giorgi and J. P. Perdew for posing the question of how to obtain the 1-matrix by contracting the 2-matrix in the case of the spin-polarized HEG. One of the authors (P.Z.) gratefully acknowledges P. Fulde for supporting this work. The other author (F.T.) thanks H. Eschrig for his support of this work and the Technische Universität Dresden for a scholarship.

Appendix 1: The case of 'no interaction'

Because of $T \sim 1/r_s^2$ and $V \sim 1/r_s$, this case is realized for $r_s \to 0$, what effects the following simplifications, where $N_\uparrow/N = (1+\zeta)/2$, $N_\downarrow/N = (1-\zeta)/2$ and $k_{\mathrm{F}\uparrow}/k_\mathrm{F} = (1+\zeta)^{1/3}$, $k_{\mathrm{F}\downarrow}/k_\mathrm{F} = (1-\zeta)^{1/3}$, and wavelengths are measured in units of k_F, lengths in units of k_F^{-1}, and energies in a.u.:

$$n_\uparrow^0(k) = \theta((1+\zeta)^{1/3} - k), \quad n_\downarrow^0(k) = \theta((1-\zeta)^{1/3} - k), \tag{60}$$

$$f_\uparrow^0(r) = 3\frac{j_1((1+\zeta)^{1/3}r)}{(1+\zeta)^{1/3}r}, \quad f_\downarrow^0(r) = 3\frac{j_1((1-\zeta)^{1/3}r)}{(1-\zeta)^{1/3}r}, \tag{61}$$

$$\frac{t_\uparrow^0}{t^0} = \frac{1}{2}(1+\zeta)^{2/3}, \quad \frac{t_\downarrow^0}{t^0} = \frac{1}{2}(1-\zeta)^{2/3}, \quad t^0 = \frac{3}{10}\frac{1}{\alpha_0^2}, \quad \alpha_0 = \left(\frac{4}{9\pi}\right)^{1/3} r_s. \tag{62}$$

There are two Fermi spheres, a large one for spin ↑ and a smaller one for spin ↓. There are three regions of different occupancies: ↑↓ in the inner sphere, ↑ in the shell, and 0 outside the shell. Because of the idempotency $[n_s^0(k)]^2 = n_s^0(k)$ and the projection $n_\uparrow^0(k)n_\downarrow^0(k) = n_\downarrow^0(k)$ from Eq.(19) it follows

$$\begin{aligned}\beta_\uparrow^0(r) = \frac{1+\zeta}{2} f_\uparrow^0(r), \quad & \beta_\downarrow^0(r) = \beta_a^0(r) = \frac{1-\zeta}{2} f_\uparrow^0(r), \\ b_\uparrow^0 = \frac{1+\zeta}{2}, \qquad & b_\downarrow^0 = b_a^0 = \frac{1-\zeta}{2}.\end{aligned} \tag{63}$$

All the quantities and relations of Table II hold [with the replacements HF → 0 and $f_s(r) \to f_s^0(r)$] also for the ideal Fermi gas including Eq.(63) and

$$\alpha_{3,4}^0(r) = \alpha^0(r) = \frac{1-\zeta}{2} f_\downarrow^0(r), \quad \tilde{\alpha}^0(r) = 0. \tag{64}$$

It follows $a^0 = (1-\zeta)/2$, what can be proved also in the following way. Starting with the definition $a = -\mathrm{Tr}\gamma_a^{3,4}/N$ one has to evaluate $\mathrm{Tr}\gamma_a^3$ or $\mathrm{Tr}\gamma_a^4$ for 'no interaction', which allows with Eq.(60) to calculate the traces in the $\mathbf{k}$ space:

$$\mathrm{Tr}\gamma_a^3 = \sum_{\mathbf{k}_{1,2}} \langle \hat{a}^\dagger_{\downarrow\mathbf{k}_1} \hat{a}^\dagger_{\uparrow\mathbf{k}_2} \hat{a}_{\downarrow\mathbf{k}_2} \hat{a}_{\uparrow\mathbf{k}_1} \rangle = \sum_{\mathbf{k}_{1,2}} \langle \hat{a}^\dagger_{\downarrow\mathbf{k}_1} \hat{a}_{\uparrow\mathbf{k}_1} \hat{a}^\dagger_{\uparrow\mathbf{k}_2} \hat{a}_{\downarrow\mathbf{k}_2} \rangle - \sum_{\mathbf{k}} \langle \hat{a}^\dagger_{\downarrow\mathbf{k}} \hat{a}_{\uparrow\mathbf{k}} \rangle = -N_\downarrow. \tag{65}$$

Thereby $\hat{a}^\dagger_{\uparrow\mathbf{k}_2} \hat{a}_{\downarrow\mathbf{k}_2}\rangle$ should be a state with a hole at $\mathbf{k}_2$ in the ↓-sphere (hence $k_2 < k_{F\downarrow}$) and an additional particle at $\mathbf{k}_2$ in the ↑-sphere. This is not possible: $\left(\hat{a}^\dagger_{\uparrow\mathbf{k}}\right)^2 = 0$. From $\mathrm{Tr}\gamma_a^4$ follows the same result:

$$\mathrm{Tr}\gamma_a^4 = \sum_{\mathbf{k}_{1,2}} \langle \hat{a}^\dagger_{\uparrow\mathbf{k}_1} \hat{a}^\dagger_{\downarrow\mathbf{k}_2} \hat{a}_{\uparrow\mathbf{k}_2} \hat{a}_{\downarrow\mathbf{k}_1} \rangle = \textstyle\sum_{\mathbf{k}_{1,2}} \langle \hat{a}^\dagger_{\uparrow\mathbf{k}_1} \hat{a}_{\downarrow\mathbf{k}_1} \hat{a}^\dagger_{\downarrow\mathbf{k}_2} \hat{a}_{\uparrow\mathbf{k}_2} \rangle - \sum_{\mathbf{k}} \langle \hat{a}^\dagger_{\uparrow\mathbf{k}} \hat{a}_{\downarrow\mathbf{k}} \rangle. \tag{66}$$

In the first term, with $\hat{a}_{\uparrow\mathbf{k}_2}$ a hole in the ↑-sphere can be made only in the shell $k_{F\downarrow} < k_2 < k_{F\uparrow}$ between the two Fermi surfaces. Because k_2 is thus outside the ↓-sphere, the ↓ places in the shell are empty and can be occupied: $\hat{a}^\dagger_{\downarrow\mathbf{k}_2} \hat{a}_{\uparrow\mathbf{k}_2}\rangle \neq 0$. With $\mathbf{k}_1 = \mathbf{k}_2$ and $\sum_{k_{F\downarrow}<k<k_{F\uparrow}} 1 = N_\uparrow - N_\downarrow$, again $a^0 = N_\downarrow/N = (1-\zeta)/2$ turns out.

The PDs for spin-parallel pairs are

$$\begin{aligned} g_{\uparrow\uparrow}^0(r) = 1 - |f_\uparrow^0(r)|^2, \quad & \int d^3r\, \rho[1 - g_{\uparrow\uparrow}^0(r)] = \frac{2}{1+\zeta}, \quad g_{\uparrow\uparrow}^0(\infty) = 1, \\ g_{\downarrow\downarrow}^0(r) = 1 - |f_\downarrow^0(r)|^2, \quad & \int d^3r\, \rho[1 - g_{\downarrow\downarrow}^0(r)] = \frac{2}{1-\zeta}, \quad g_{\downarrow\downarrow}^0(\infty) = 1. \end{aligned} \tag{67}$$

The PDs for the spin-antiparallel pairs are

$$\begin{aligned} g_a^0(r) = 1, \qquad & \textstyle\int d^3r\, \rho[1 - g_a^0(r)] = 0, \qquad & g_a^0(\infty) = 1, \\ g_a'^0(r) = -\mathrm{Re} f_\uparrow^0(r) f_\downarrow^0(r), \qquad & \textstyle\int d^3r\, \rho g_a'^0(r) = \frac{2}{1+\zeta}, \qquad & g_a'^0(\infty) = 0, \\ g_\pm^0(r) = 1 \mp \mathrm{Re} f_\uparrow^0(r) f_\downarrow^0(r), \qquad & \textstyle\int d^3r\, \rho[1 - g_\pm^0(r)] = \mp\frac{2}{1+\zeta}, \qquad & g_\pm(\infty) = 1. \end{aligned} \tag{68}$$

Finally the spin-weighted PD is

$$\begin{aligned} g^0(r) &= \left(\frac{1+\zeta}{2}\right)^2 g^0_{\uparrow\uparrow} + \left(\frac{1-\zeta}{2}\right)^2 g^0_{\downarrow\downarrow} + \frac{1-\zeta^2}{2} \quad \text{or} \\ 1 - g^0(r) &= \left(\frac{1+\zeta}{2}\right)^2 |f^0_\uparrow(r)|^2 + \left(\frac{1-\zeta}{2}\right)^2 |f^0_\downarrow(r)|^2 \end{aligned} \tag{69}$$

with $\int d^3r\ \rho[1 - g^0(r)] = 1, \quad g^0(\infty) = 1$, besides $\Delta g^0(r) = g_a'^0(r)$.

Appendix 2: The case of 'no spin-polarization'

The simplifications for 'no polarization' are: $\zeta = 0$, consequently $N_\uparrow = N_\downarrow = N/2$, and also $\rho_\uparrow = \rho_\downarrow = \rho/2$, $\gamma_\uparrow = \gamma_\downarrow = \gamma/2$, $f_\uparrow = f_\downarrow = f$, $\gamma = \rho f$, $n_\uparrow = n_\downarrow = n$. In this case the spin and position matrices are

$$\begin{aligned} &\delta_p \equiv \delta_{\uparrow\uparrow} + \delta_{\downarrow\downarrow} = \delta_{\sigma_1\sigma_1'}\delta_{\sigma_2\sigma_2'}\delta_{\sigma_1\sigma_2}, \quad \langle\uparrow\uparrow\uparrow\uparrow\rangle = \langle\downarrow\downarrow\downarrow\downarrow\rangle \quad \text{or} \quad \gamma_{\uparrow\uparrow} = \gamma_{\downarrow\downarrow} \equiv \gamma_p, \ \ \mathrm{Tr}\gamma_p = \frac{N}{2}\left(\frac{N}{2} - 1\right), \\ &\delta_a \equiv \delta_a^1 + \delta_a^2 = \delta_{\sigma_1\sigma_1'}\delta_{\sigma_2\sigma_2'}\delta_{\sigma_1,-\sigma_2}, \quad \langle\uparrow\downarrow\downarrow\uparrow\rangle = \langle\downarrow\uparrow\uparrow\downarrow\rangle \quad \text{or} \quad \gamma_a^1 = \gamma_a^2 \equiv \gamma_a, \ \ \mathrm{Tr}\gamma_a = \left(\frac{N}{2}\right)^2, \\ &\delta_a' \equiv \delta_a^3 + \delta_a^4 = \delta_{\sigma_1\sigma_2'}\delta_{\sigma_2\sigma_1'}\delta_{\sigma_1,-\sigma_2}, \quad \langle\downarrow\uparrow\downarrow\uparrow\rangle = \langle\uparrow\downarrow\uparrow\downarrow\rangle \quad \text{or} \quad \gamma_a^3 = \gamma_a^4 \equiv \gamma_a', \ \ \mathrm{Tr}\gamma_a' = -\frac{N}{2}. \end{aligned} \tag{70}$$

An immediate consequence is $\tilde{\gamma}_a = \tilde{\gamma}_a' = \tilde{\gamma}_\pm = 0$. Besides from Table I it follows $\gamma_a^{\mathrm{HF}} + \gamma_a'^{\mathrm{HF}} = \gamma_p^{\mathrm{HF}}$ and the Feynman diagrams (42) and (44) lead correspondingly to $\chi_a + \chi_a' = \chi_p$, because the spin-directions ↑ and ↓ are equally weighted. (Note also, that γ_p and $\gamma_- = \gamma_a + \gamma_a'$ have the same symmetry property, namely changing sign under the replacements $\mathbf{r}_1 \leftrightarrow \mathbf{r}_2$ or $\mathbf{r}_1' \leftrightarrow \mathbf{r}_2'$.) So it is

$$\gamma_a + \gamma_a' = \gamma_p \quad \text{or} \quad \gamma_a' = \gamma_p - \gamma_a, \quad \text{or} \quad \gamma_- = \gamma_p \tag{71}$$

(for $r_s = 0$ this can be proved directly). Eq. (71) says that for $\zeta = 0$ the 2-matrix γ_2 consists of only two components, namely γ_p and γ_a or γ_+ and γ_-. Indeed the parallel/antiparallel representation (24) takes the simple form

$$\begin{aligned} \gamma_2 &= (\delta_p + \delta_a')\gamma_p + (\delta_a - \delta_a')\gamma_a, \\ \mathrm{Tr}\gamma_2 &= 2 \cdot \frac{N}{2}\left(\frac{N}{2} - 1\right) + 2 \cdot \left(\frac{N}{2}\right)^2 = N(N-1) \end{aligned} \tag{72}$$

and the equivalent triplet/singlet representation (25) simplifies as

$$\begin{aligned} \gamma_2 &= (\delta_p + \delta_+)\gamma_- + \delta_-\gamma_+ \\ \mathrm{Tr}\gamma_2 &= 3 \cdot \frac{N}{2}\left(\frac{N}{2} - 1\right) + 1 \cdot \frac{N}{2}\left(\frac{N}{2} + 1\right) = N(N-1). \end{aligned} \tag{73}$$

Its equivalence with Eq.(72) follows from Eq.(71) and $\gamma_a = (\gamma_+ + \gamma_-)/2$ and $\delta_\pm = (\delta_a \pm \delta'_a)/2$ of Table I. The spin structure of Eq.(73) agrees with the result of the RPA-like treatment for small r_s [38]. Note $\mathrm{Tr}\gamma_\pm = \frac{N}{2}(\frac{N}{2} \pm 1)$. The spin-matrices appearing in Eqs.(72), (73) are

$$\begin{aligned}
\delta_p + \delta'_a &= \delta_{\sigma_1,\sigma'_1}\delta_{\sigma_2,\sigma'_2}\delta_{\sigma_1,\sigma_2} + \delta_{\sigma_1,\sigma'_2}\delta_{\sigma_2,\sigma'_1}\delta_{\sigma_1,-\sigma_2} &&\to \delta_{\sigma_1,\sigma_2}, \\
\delta_p + \delta_+ &= \tfrac{1}{2}(\delta_{\sigma_1,\sigma'_1}\delta_{\sigma_2,\sigma'_2} + \delta_{\sigma_1,\sigma'_2}\delta_{\sigma_2,\sigma'_1}) &&\to \frac{1}{2}(1 + \delta_{\sigma_1,\sigma_2}), \\
\delta_- = \frac{1}{2}(\delta_a - \delta'_a) &= \tfrac{1}{2}(\delta_{\sigma_1,\sigma'_1}\delta_{\sigma_2,\sigma'_2} - \delta_{\sigma_1,\sigma'_2}\delta_{\sigma_2,\sigma'_1}) &&\to \frac{1}{2}\delta_{\sigma_1,-\sigma_2}.
\end{aligned} \tag{74}$$

(In the last line it holds $\frac{1}{2}(\cdots) = \frac{1}{2}(\cdots)\delta_{\sigma_1,-\sigma_2}$, because of $1 = \delta_{\sigma_1,\sigma_2} + \delta_{\sigma_1,-\sigma_2}$ and $\frac{1}{2}(\cdots)\delta_{\sigma_1,\sigma_2} = 0$.) The arrows indicate the diagonal elements, which give the spin-dependent PD $\mathrm{D}\gamma_2 = \rho_2 = \rho^2 g$ and the spin-summed PD $g(r) = \sum\limits_{\sigma_{1,2}} g(1,2)$ equivalently as

$$\begin{aligned}
g(1,2) &= \tfrac{1}{4}[\delta_{\sigma_1,\sigma_2}g_p(r_{12}) + \delta_{\sigma_1,-\sigma_2}g_a(r_{12})], & g(r) &= \frac{1}{2}[g_p(r) + g_a(r)] \quad \text{or} \\
g(1,2) &= \tfrac{1}{4}\left[\tfrac{1}{2}(1 + \delta_{\sigma_1,\sigma_2})g_-(r_{12}) + \tfrac{1}{2}\delta_{\sigma_1,-\sigma_2}g_+(r_{12})\right], & g(r) &= \frac{1}{4}[3g_-(r) + g_+(r)],
\end{aligned} \tag{75}$$

in agreement with Eq.(31). From the above relations between the p, a and the ± components of the 2-matrix follow corresponding relations for the PD:

$$\begin{aligned}
\gamma_p = \gamma_-, \gamma = \frac{1}{2}(\gamma_+ + \gamma_-) \quad &\text{or} \quad \gamma_+ = 2\gamma_a - \gamma_p, \gamma_- = \gamma_p, \\
g_p = g_-, g_a = \frac{1}{2}(g_+ + g_-) \quad &\text{or} \quad g_+ = 2g_a - g_p, g_- = g_p.
\end{aligned} \tag{76}$$

Whereas in general $\Delta g = (g_+ - g_-)/2$ (= singlet-triplet PD splitting), for $\zeta = 0$ it is $\Delta g = g_a - g_p$ (= antiparallel-parallel PD difference) or $g_p = g - \Delta g/2$, $g_a = g + \Delta g/2$. So, Δg may be addressed as magnetic PD. The contraction of Eq.(71) yields $\frac{\rho}{2}f(\frac{N}{2} - 1) = \frac{\rho}{2}f\frac{N}{2} - \rho\alpha$ or $\alpha(r) = f(r)/2$, thus for $r = 0$ it follows $a = 1/2$. In the contraction and normalization of the HF and cumulant parts appear $\beta_s = \beta_a \equiv \beta$, thus $b_s = b_a \equiv b$ and $c_s = c_a \equiv c$, $b = (1 - b)/2$. The case $\zeta = 0$ is summarized in Table IV.

The PDs are defined and normalized according to (with $\mathbf{r}'_1 = \mathbf{r}_1$, $\mathbf{r}'_2 = \mathbf{r}_2$, $r = r_{12}$)

$$\left(\frac{\rho}{2}\right)^2 g_\pm(r) = \gamma_\pm(\mathbf{r}_1|\mathbf{r}_1, \mathbf{r}_2|\mathbf{r}_2) \ , \quad \int d^3r\ \rho[1 - g_\pm(r)] = \mp 2,\ g_\pm(\infty) = 1. \tag{77}$$

With the functions

$$\left(\frac{\rho}{2}\right)^2 G_\pm(r_{11'}, r_{12}) = \int \frac{d\Omega_{12}}{4\pi}\gamma_\pm(\mathbf{r}_1|\mathbf{r}'_1, \mathbf{r}_2|\mathbf{r}_2) \tag{78}$$

the contraction SRs are

$$\int d^3r_{12}\ \rho[f(r_{11'}) - G_\pm(r_{11'}, r_{12})] = \mp 2f(r_{11'})\ ,\quad G_\pm(r_{11'}, \infty) = f(r_{11'}). \tag{79}$$

From the spectral resolution (54) it follows

$$\begin{aligned} G_\pm(r_{11'}, r_{12}) &= \int \frac{d\Omega_{12}}{4\pi}\frac{4}{\rho^2}\sum_{\mathbf{K},\mathbf{k}} n(|\frac{1}{2}\mathbf{K}+\mathbf{k}|)n(|\frac{1}{2}\mathbf{K}-\mathbf{k}|)\varphi_\pm(\mathbf{r}_1, \mathbf{r}_2)\varphi_\pm^*(\mathbf{r}_1', \mathbf{r}_2), \\ \varphi_\pm(\mathbf{r}_1, \mathbf{r}_2) &= \frac{1}{\sqrt{\Omega}}\mathrm{e}^{\mathrm{i}\mathbf{KR}}\frac{4\pi}{\sqrt{\Omega}}\sum_L^{\pm} \mathrm{i}^l R_L(\mathbf{r}, \mathbf{k}), \quad R_L(\mathbf{r}, \mathbf{k}) = R_l(r, k)Y_L(\mathbf{e}_r)Y_L^*(\mathbf{e}_k). \end{aligned} \tag{80}$$

Here the functions $G_\pm$ contain the PDs $g_\pm(r) = G_\pm(0, r)$, and the 1-matrix $f(r) = G_\pm(r, \infty)$. For 'no polarization' there are the contraction SRs (79) and $f(0) = 1$.

Appendix 3. The case of 'full spin polarization'

Here $\zeta = 1$ is considered, i.e. $N_\downarrow = 0$, thus $\rho_\downarrow, \gamma_\downarrow, f_\downarrow, n_\downarrow = 0$ and $N_\uparrow = N$. The above formulae simplify as

$$\gamma_1(1|1') = \delta_{\sigma_1,\sigma_1'}\delta_{\sigma_1,\uparrow}\gamma(\mathbf{r}_1|\mathbf{r}_1'), \gamma(\mathbf{r}_1|\mathbf{r}_1') = \rho f(r_{11'}), f(0) = 1, \tag{81}$$

$$\gamma_2(1|1', 2|2') = \delta_+(\sigma_1|\sigma_1', \sigma_2|\sigma_2')\delta_{\sigma_1,\uparrow}\delta_{\sigma_2,\uparrow}\gamma(\mathbf{r}_1|\mathbf{r}_1', \mathbf{r}_2|\mathbf{r}_2'), \quad \mathrm{Tr}\gamma_2 = N(N-1), \tag{82}$$

$$\rho^2 g(r_{12}) = \gamma(\mathbf{r}_1|\mathbf{r}_1, \mathbf{r}_2|\mathbf{r}_2), \int d^3r \rho[1 - g(r)] = 1, \quad g(\infty) = 1, \tag{83}$$

$$\rho^2 G(r_{11'}, r_{12}) = \int \frac{d\Omega_{12}}{4\pi}\gamma(\mathbf{r}_1|\mathbf{r}_1', \mathbf{r}_2|\mathbf{r}_2), \quad G(0, r_{12}) = g(r_{12}), \tag{84}$$

$$\int d^3r_{12}\ \rho[f(r_{11'}) - G(r_{11'}, r_{12})] = f(r_{11'}), \quad G(r_{11'}, \infty) = f(r_{11'}). \tag{85}$$

Note $g_{\downarrow\downarrow}(r) = 0$, thus $\int d^3r\ \rho[1 - g_{\downarrow\downarrow}(r)] = 2/(1-\zeta)$ diverges for $\zeta \to 1$.

Appendix 4: The spin structure of γ_2

Eq.(25) agrees term by term with the Eq.(6-13) of Ref. [12], where the block-diagonality of γ_2 in the representation corresponding to the common eigenstates of the 2-body spin-operators $\hat{s}_{z1} + \hat{s}_{z2}$ and $(\hat{\mathbf{s}}_1 + \hat{\mathbf{s}}_2)^2$:

$$\gamma_2 = (C_{11}D_{111}C_{11}' + C_{1-1}D_{11-1}C_{1-1}' + C_{10}D_{110}C_{10}') + C_{00}D_{000}C_{00}' + (C_{10}D_{100}C_{00}' + C_{00}D_{010}C_{10}') \tag{86}$$

Here C_{SM}, C'_{SM}, $D_{SS'M}$ are abbreviations for $C^{\sigma_1\sigma_2}_{SM}$, $C^{\sigma'_1\sigma'_2}_{SM}$, $D_{SS'M}(\mathbf{r}_1, \mathbf{r}_2|\mathbf{r}'_1, \mathbf{r}'_2)$, respectively. For the position matrices it is e.g. $D_{111} = \gamma_{\uparrow\uparrow}$, $D_{11-1} = \gamma_{\downarrow\downarrow}$, $D_{110} = \gamma_{a-}$. The triplet/singlet spin functions or (Clebsch-Gordan coefficients)

$$\begin{aligned} C_{11} &= \delta_{\sigma_1\uparrow}\delta_{\sigma_2\uparrow}, & C_{1-1} &= \delta_{\sigma_1\downarrow}\delta_{\sigma_2\downarrow}, \\ C_{10} &= \tfrac{1}{\sqrt{2}}(\delta_{\sigma_1\uparrow}\delta_{\sigma_2\downarrow} + \delta_{\sigma_2\uparrow}\delta_{\sigma_1\downarrow}), & C_{00} &= \frac{1}{\sqrt{2}}(\delta_{\sigma_1\uparrow}\delta_{\sigma_2\downarrow} - \delta_{\sigma_2\uparrow}\delta_{\sigma_1\downarrow}) \end{aligned} \tag{87}$$

give the corresponding spin matrices of Eqs.(22) and (26) according to

$$\begin{aligned} \delta_{\uparrow\uparrow} &= C_{11}C'_{11}, & \delta_{\downarrow\downarrow} &= C_{1-1}C'_{1-1}, & \delta_+ &= C_{10}C'_{10}, \\ \delta_- &= C_{00}C'_{00}, & \tilde{\delta}_+ &= C_{10}C'_{00}, & \tilde{\delta}_- &= C_{00}C'_{10}. \end{aligned} \tag{88}$$

Table I. The 2-matrices of the spin-polarized HEG and their properties (cf. Sec. 5). The δ_{ss} and δ^1_a etc. are defined in Eq.(22). The operators D, P, Q, C, and Tr are defined after Eq.(21). The lines $\uparrow\uparrow$ and $\downarrow\downarrow$ are for spin-parallel electron pairs, the following lines for spin-antiparrallel electron pairs. The 2-matrices depend on $\mathbf{r}_1|\mathbf{r}'_1, \mathbf{r}_2|\mathbf{r}'_2$, the PDs of column D$\gamma$ depend on $r_{12} = |\mathbf{r}_1 - \mathbf{r}_2|$. In column C$\gamma$ the 1-matrices $f_\uparrow$, $f_\downarrow$, $\alpha_{3,4}$ as well as $f = \frac{1}{2}(f_\uparrow + f_\downarrow)$, $\tilde{f} = \frac{1}{2}(f_\uparrow - f_\downarrow)$, $\alpha = \frac{1}{2}(\alpha_3 + \alpha_4)$, $\tilde{\alpha} = \frac{1}{2}(\alpha_3 - \alpha_4)$ appear, they depend on $r_{11'} = |\mathbf{r}_1 - \mathbf{r}'_1|$. Note $f_\uparrow(0) = f_\downarrow(0) = f(0) = 1$ and $\alpha_3(0) = \alpha_4(0) = \alpha(0) = a$ (which is a function of r_s and ζ), hence $\tilde{f}(0) = 0$ and $\tilde{\alpha}(0) = 0$.

δ	Trδ	γ	Dγ	Pγ	Qγ	Cγ	Tr γ
$\delta_{\uparrow\uparrow}$	1	$\gamma_{\uparrow\uparrow} = \langle\uparrow\uparrow\uparrow\uparrow\rangle$	$\rho_\uparrow^2 g_{\uparrow\uparrow}$	$-\gamma_{\uparrow\uparrow}$	$\gamma_{\uparrow\uparrow}$	$\rho_\uparrow f_\uparrow (N_\uparrow - 1)$	$N_\uparrow (N_\uparrow - 1)$
$\delta_{\downarrow\downarrow}$	1	$\gamma_{\downarrow\downarrow} = \langle\downarrow\downarrow\downarrow\downarrow\rangle$	$\rho_\downarrow^2 g_{\downarrow\downarrow}$	$-\gamma_{\downarrow\downarrow}$	$\gamma_{\downarrow\downarrow}$	$\rho_\downarrow f_\downarrow (N_\downarrow - 1)$	$N_\downarrow (N_\downarrow - 1)$
δ_a^1	1	$\gamma_a^1 = \langle\uparrow\downarrow\downarrow\uparrow\rangle$	$\rho_\uparrow \rho_\downarrow g_a$	$-\gamma_a^4$	γ_a^1	$\rho_\uparrow f_\uparrow N_\downarrow$	$N_\uparrow N_\downarrow$
δ_a^2	1	$\gamma_a^2 = \langle\downarrow\uparrow\uparrow\downarrow\rangle$	$\rho_\downarrow \rho_\uparrow g_a$	$-\gamma_a^3$	γ_a^2	$\rho_\downarrow f_\downarrow N_\uparrow$	$N_\downarrow N_\uparrow$
δ_a^3	0	$\gamma_a^3 = \langle\downarrow\uparrow\downarrow\uparrow\rangle$	$-\rho_\uparrow \rho_\downarrow \Delta g$	$-\gamma_a^2$	γ_a^4	$-\rho\alpha_3$	$-Na$
δ_a^4	0	$\gamma_a^4 = \langle\uparrow\downarrow\uparrow\downarrow\rangle$	$-\rho_\downarrow \rho_\uparrow \Delta g$	$-\gamma_a^1$	γ_a^3	$-\rho\alpha_4$	$-Na$
$\delta_a = \delta_a^1 + \delta_a^2$	2	$\gamma_a = \frac{1}{2}(\gamma_a^1 + \gamma_a^2)$	$\rho_\uparrow \rho_\downarrow g_a$	$-\gamma_a'$	γ_a	$\rho_\uparrow f N_\downarrow$	$N_\uparrow N_\downarrow$
$\delta_a' = \delta_a^3 + \delta_a^4$	0	$\gamma_a' = \frac{1}{2}(\gamma_a^3 + \gamma_a^4)$	$-\rho_\downarrow \rho_\uparrow \Delta g$	$-\gamma_a$	γ_a'	$-\rho\alpha$	$-Na$
$\tilde{\delta}_a = \delta_a^1 - \delta_a^2$	0	$\tilde{\gamma}_a = \frac{1}{2}(\gamma_a^1 - \gamma_a^2)$	0	$+\tilde{\gamma}_a'$	$\tilde{\gamma}_a$	$\rho_\downarrow \tilde{f} N_\uparrow$	0
$\tilde{\delta}_a' = \delta_a^3 - \delta_a^4$	0	$\tilde{\gamma}_a' = \frac{1}{2}(\gamma_a^3 - \gamma_a^4)$	0	$+\tilde{\gamma}_a$	$-\tilde{\gamma}_a'$	$-\rho\tilde{\alpha}$	0
$\delta_- = \frac{1}{2}(\delta_a - \delta_a')$	1	$\gamma_+ = \gamma_a - \gamma_a'$	$\rho_\uparrow \rho_\downarrow (g_a + \Delta g)$	$+\gamma_+$	γ_+	$\rho_\uparrow f N_\downarrow + \rho\alpha$	$N_\uparrow N_\downarrow + Na$
$\delta_+ = \frac{1}{2}(\delta_a + \delta_a')$	1	$\gamma_- = \gamma_a + \gamma_a'$	$\rho_\downarrow \rho_\uparrow (g_a - \Delta g)$	$-\gamma_-$	γ_-	$\rho_\downarrow f N_\uparrow - \rho\alpha$	$N_\downarrow N_\uparrow - Na$
$\tilde{\delta}_- = \frac{1}{2}(\tilde{\delta}_a + \tilde{\delta}_a')$	0	$\tilde{\gamma}_+ = \tilde{\gamma}_a + \tilde{\gamma}_a'$	0	$+\tilde{\gamma}_+$	$\tilde{\gamma}_-$	$\rho_\uparrow \tilde{f} N_\downarrow - \rho\tilde{\alpha}$	0
$\tilde{\delta}_+ = \frac{1}{2}(\tilde{\delta}_a - \tilde{\delta}_a')$	0	$\tilde{\gamma}_- = \tilde{\gamma}_a - \tilde{\gamma}_a'$	0	$-\tilde{\gamma}_-$	$\tilde{\gamma}_+$	$\rho_\downarrow \tilde{f} N_\uparrow + \rho\tilde{\alpha}$	0

Table II. The HF parts of the cumulant expansion (cf. Sec. 8). Note $\alpha_{3,4}^{\mathrm{HF}}(r) = \alpha^{\mathrm{HF}}(r) = \beta_a^{\mathrm{HF}}(\mathrm{r})$, $a^{\mathrm{HF}} = b_a^{\mathrm{HF}}$, $\tilde{\alpha}^{\mathrm{HF}} = 0$. For 'no interaction' the cumulant parts vanish, only the HF parts remain, which simplify furthermore as $\beta_s^0(r) = \frac{N_s}{N} f_s^0(\mathrm{r})$, $\beta_a^0(r) = \frac{N_\downarrow}{N} f_\downarrow^0(r)$, thus $b_s^0 = \frac{N_s}{N}$, $b_a^0 = \frac{N_\downarrow}{N}$.

γ^{HF}	$\mathrm{C}\gamma^{\mathrm{HF}}$	$\mathrm{Tr}\gamma^{\mathrm{HF}}$
$\gamma_{ss}^{\mathrm{HF}} = \rho_s^2[f_s(r_{11'})f_s(r_{22'}) - f_s(r_{12'})f_s(r_{21'})]$	$\rho_s f_s N_s - \rho\beta_s$	$N_s^2 - Nb_s$
$\gamma_a^{1\mathrm{HF}} = \rho_\uparrow\rho_\downarrow f_\uparrow(r_{11'})f_\downarrow(r_{22'})$	$\rho_\uparrow f_\uparrow N_\downarrow$	$N_\uparrow N_\downarrow$
$\gamma_a^{2\mathrm{HF}} = \rho_\downarrow\rho_\uparrow f_\downarrow(r_{11'})f_\uparrow(r_{22'})$	$\rho_\downarrow f_\downarrow N_\uparrow$	$N_\downarrow N_\uparrow$
$\gamma_a^{3\mathrm{HF}} = -\rho_\uparrow\rho_\downarrow f_\uparrow(r_{12'})f_\downarrow(r_{21'})$	$-\rho\beta_a$	$-Nb_a$
$\gamma_a^{4\mathrm{HF}} = -\rho_\downarrow\rho_\uparrow f_\downarrow(r_{12'})f_\uparrow(r_{21'})$	$-\rho\beta_a$	$-Nb_a$
$\gamma_a^{\mathrm{HF}} = \frac{1}{2}\rho_\uparrow\rho_\downarrow[f_\uparrow(r_{11'})f_\downarrow(r_{22'}) + f_\downarrow(r_{11'})f_\uparrow(r_{22'})]$	$\rho_\uparrow f N_\downarrow$	$N_\uparrow N_\downarrow$
$\gamma_a'^{\,\mathrm{HF}} = -\frac{1}{2}\rho_\uparrow\rho_\downarrow[f_\uparrow(r_{12'})f_\downarrow(r_{21'}) + f_\downarrow(r_{12'})f_\uparrow(r_{21'})]$	$-\rho\beta_a$	$-Nb_a$
$\tilde{\gamma}_a^{\,\mathrm{HF}} = \frac{1}{2}\rho_\downarrow\rho_\uparrow[f_\uparrow(r_{11'})f_\downarrow(r_{22'}) - f_\downarrow(r_{11'})f_\uparrow(r_{22'})]$	$\rho_\downarrow \tilde{f} N_\uparrow$	0
$\tilde{\gamma}_a'^{\,\mathrm{HF}} = -\frac{1}{2}\rho_\uparrow\rho_\downarrow[f_\uparrow(r_{12'})f_\downarrow(r_{21'}) - f_\downarrow(r_{12'})f_\uparrow(r_{21'})]$	0	0
$\gamma_+^{\mathrm{HF}} = \gamma_a^{\mathrm{HF}} - \gamma_a'^{\,\mathrm{HF}}$	$\rho_\uparrow f N_\downarrow + \rho\beta_a$	$N_\uparrow N_\downarrow + Nb_a$
$\gamma_-^{\mathrm{HF}} = \gamma_a^{\mathrm{HF}} + \gamma_a'^{\,\mathrm{HF}}$	$\rho_\downarrow f N_\uparrow - \rho\beta_a$	$N_\downarrow N_\uparrow - Nb_a$
$\tilde{\gamma}_+^{\mathrm{HF}} = \tilde{\gamma}_a^{\,\mathrm{HF}} + \tilde{\gamma}_a'^{\,\mathrm{HF}}$	$\rho_\uparrow \tilde{f} N_\downarrow$	0
$\tilde{\gamma}_-^{\mathrm{HF}} = \tilde{\gamma}_a^{\mathrm{HF}} - \tilde{\gamma}_a'^{\,\mathrm{HF}}$	$\rho_\downarrow \tilde{f} N_\uparrow$	0

Table III. The cumulant 2-matrices, being size-extensively normalized (cf. Sec. 8). The index ‘c’ means ‘connected’ or ‘cumulant’. The cumulant normalization $\mathrm{Tr}\chi_{ss}/N_s = c_s$ measures the correlation strength, cf. Eq.(20).

$\chi = \gamma^{\mathrm{HF}} - \gamma$	Cχ	Trχ
$\chi_{ss} = -\langle ssss \rangle_{\mathrm{c}}$	$\rho_s f_s - \rho \beta_s$	$N_s - N b_s$
$\chi_a^1 = -\langle \uparrow\downarrow\downarrow\uparrow \rangle_{\mathrm{c}}$	0	0
$\chi_a^2 = -\langle \downarrow\uparrow\uparrow\downarrow \rangle_{\mathrm{c}}$	0	0
$\chi_a^3 = -\langle \downarrow\uparrow\downarrow\uparrow \rangle_{\mathrm{c}}$	$\rho(\alpha_3 - \beta_a)$	$N(a - b_a)$
$\chi_a^4 = -\langle \uparrow\downarrow\uparrow\downarrow \rangle_{\mathrm{c}}$	$\rho(\alpha_4 - \beta_a)$	$N(a - b_a)$
$\chi_a = \frac{1}{2}(\chi_a^1 + \chi_a^2)$	0	0
$\chi'_a = \frac{1}{2}(\chi_a^3 + \chi_a^4)$	$\rho(\alpha - \beta_a)$	$N(a - b_a)$
$\tilde{\chi}_a = \frac{1}{2}(\chi_a^1 - \chi_a^2)$	0	0
$\tilde{\chi}'_a = \frac{1}{2}(\chi_a^3 - \chi_a^4)$	$\rho\tilde{\alpha}$	0
$\chi_+ = \chi_a - \chi'_a$	$-\rho(\alpha - \beta_a)$	$-N(a - b_a)$
$\chi_- = \chi_a + \chi'_a$	$+\rho(\alpha - \beta_a)$	$+N(a - b_a)$
$\tilde{\chi}_+ = \tilde{\chi}_a + \tilde{\chi}'_a$	$+\rho\tilde{\alpha}$	0
$\tilde{\chi}_- = \tilde{\chi}_a - \tilde{\chi}'_a$	$-\rho\tilde{\alpha}$	0

Table IV. This is Table I, simplified for the case of the spin-unpolarized HEG. In App. 2 it is shown : $\gamma_p = \gamma_- = \gamma_a - \gamma'_a$, thus also $\Delta g = g_a - g_p$, and $\alpha = f/2$, $a = 1/2$. Besides, the HF parts contain $\beta_s = \beta_a \equiv \beta$, thus $b_s = b_a \equiv b$, cf. Eqs.(19), (20).

γ	Dγ	Pγ	Qγ	Cγ	Trγ
$\gamma_p = \langle \uparrow\uparrow\uparrow\uparrow \rangle$	$(\frac{\rho}{2})^2 g_p$	$-\gamma_p$	γ_p	$\frac{\rho}{2} f (\frac{N}{2} - 1)$	$\frac{N}{2}(\frac{N}{2} - 1)$
$\gamma_a = \langle \uparrow\downarrow\downarrow\uparrow \rangle$	$(\frac{\rho}{2})^2 g_a$	$-\gamma'_a$	γ_a	$\frac{\rho}{2} f \frac{N}{2}$	$(\frac{N}{2})^2$
$\gamma'_a = \langle \downarrow\uparrow\downarrow\uparrow \rangle$	$-(\frac{\rho}{2})^2 \Delta g$	$-\gamma_a$	γ'_a	$-\frac{\rho}{2} f$	$-\frac{N}{2}$
$\gamma_+ = \gamma_a - \gamma'_a$	$(\frac{\rho}{2})^2 (g_a + \Delta g)$	$+\gamma_+$	γ_+	$\frac{\rho}{2} f (\frac{N}{2} + 1)$	$\frac{N}{2}(\frac{N}{2} + 1)$
$\gamma_- = \gamma_a + \gamma'_a$	$(\frac{\rho}{2})^2 (g_a - \Delta g)$	$-\gamma_-$	γ_-	$\frac{\rho}{2} f (\frac{N}{2} - 1)$	$\frac{N}{2}(\frac{N}{2} - 1)$

[1] K. S. Singwi and M. P. Tosi, in *Solid State Physics*, ed. by H. Ehrenreich, F. Seitz,and D. Turnbull, Academic, New York, 1981, Vol. 36, p. 177; P. Fulde, *Electron Correlation in Molecules and Solids*, 3rd ed., Springer, Berlin, 1995; N. H. March, *Electron Correlation in Molecules and Condensed Phases*, Plenum, New York, 1996; N. H. March (ed.), *Electron Correlation in the Solid State*, Imperial College Press, London, 1999; W. Kutzelnigg, "Theory of Electron Correlation", in J. Rychlewski (ed.), *Explicitly Correlated Wave Functions in Chemistry and Physics. Theory and Applications*, Kluwer, Dordrecht, 2003, p. 3.

[2] A. Gonis, N. Kioussis, and M. Ciftan (Eds.), *Electron Correlations and Materials Properties*, Kluwer/Plenum, New York, 1999.

[3] A. Gonis, N. Kioussis, and M. Ciftan (Eds.), *Electron Correlations and Materials Properties* 2, Kluwer/Plenum, New York, 2002.

[4] D.M. Ceperley, Phys. Rev. B **18**, 3126 (1978); D.M. Ceperley and B.J. Alder, Phys. Rev. Lett. **45**, 566 (1980).

[5] J. P. Perdew and Y. Wang, Phys. Rev. B **46**, 12947 (1992); **56**, 7018 (1997).

[6] G. Ortiz and P. Ballone, Phys. Rev. B **50**, 1391 (1994); **56**, 9970 (1997); G. Ortiz, M. Harris, and P. Ballone, Phys. Rev. Lett. **82**, 5317 (1999).

[7] P. Gori-Giorgi, F. Sacchetti, and G.B. Bachelet, Phys. Rev. B **61**, 7353 (2000); **66**, 159901(E) (2002).

[8] P. Gori-Giorgi and J. P. Perdew, Phys. Rev. B **66**, 165118 (2002).

[9] P. Ziesche, phys. stat. sol. (b) **232**, 231 (2002); P. Gori-Giorgi and P. Ziesche, Phys. Rev. B **66**, 235116 (2002).

[10] N. H. March, Phys. Rev. **110**, 604 (1958).

[11] W. Macke and P. Ziesche, Ann. Physik (Leipzig) **13**, 25 (1964).

[12] E. R. Davidson, *Reduced Density Matrices in Quantum Chemistry*, Academic, New York, 1976.

[13] J. Cioslowski (ed.), *Many-Electron Densities and Reduced Density Matrices*, Kluwer/Plenum, New York, 2000.

[14] A. J. Coleman and V. I. Yukalov, *Reduced Density Matrices. Coulson's Challenge*, Springer, Berlin, 2000; A. J. Coleman, Phys. Rev. A **66**, 022503 (2002); A. Beste, K. Runge, and R. Bartlett, Chem. Phys. Lett. **355**, 263 (2002).

[15] E.R. Davidson, Int. J. Quantum Chem. **91**, 1 (2003) and refs. therein.

[16] C. Caccamo, G. Pizzimenti, and M. P. Tosi, Nuovo Cimento **31**B, 53 (1976).

[17] J. C. Kimball, Phys. Rev. A **7**, 1648 (1973).

[18] A. W. Overhauser, Can. J. Phys. **73**, 683 (1995).

[19] P. Gori-Giorgi and J. P. Perdew, Phys. Rev. B **64**, 155102 (2001); P. Gori-Giorgi, in Ref. [3], p. 379; M. Corona, P. Gori-Giorgi, and J.P. Perdew, unpublished.

[20] B. Davoudi, M. Polini, R. Asgari, and M.P. Tosi, Phys. Rev. B **66**, 075110 (2002); cond-mat/0206456; B. Davoudi, R. Asgari, M. Polini, and M.P. Tosi, unpublished.

[21] P. Ziesche, Phys. Rev. B **67**, 233102 (2003),

[22] P. Ziesche, K. Pernal, and F. Tasnádi, phys. stat. sol. (b) **239**, 185 (2003).

[23] P. Ziesche and F. Tasnádi, submitted to Ann. Physik (Pobierowo Proceedings, ed. by J. Cioslowski).

[24] The general parameter theorem $\partial E/\partial\lambda = \langle \partial\hat{H}/\partial\lambda\rangle$ for bound states of $\hat{H}$ is due to P. Güttinger, Z. Phys. **73**, 169 (1932), cf. his Eq.(11). It is later given by W. Pauli, in *Handbuch der Physik*, edited by H. Geiger and K. Scheel (Springer, Berlin, 1933), Band XXIV, Teil 1, p. 162; reprinted in *Handbuch der Physik*, edited by S. Flügge (Springer, Berlin, 1958), Band V, Teil 1, p. 83. λ can be any parameter (coupling constant, nuclear coordinate, ...). For a coupling constant λ and in lowest order of perturbation theory, the theorem was given already by E. Schrödinger, Ann. Physik (Leipzig) [4], **80**, 437 (1926), cf. his Eq.(7”). The theorem is also implicitly contained in Eq.(28) of the paper by M. Born and V. Fock, Z. Phys. **51**, 165 (1928). - Hellmann and later Feynman explicitly referred to the special case of nuclear coordinates within the Born-Oppenheimer approximation, leading to the ‘Hellmann-Feynman’ forces upon nuclei: H. Hellmann, *Einführung in die Quantenchemie*, Deuticke, Leipzig, 1937, pp. 61, 285 (the original russian version is of 23 October 1936: G. Gel’man, Quantenchemie, ONTI, Moscow and Leningrad, p. 428); R.P. Feynman, Phys. Rev. **56**, 340 (1939). Hellmann stressed, that the forces calculated in this way sensitively depend on the electron density used therein. For variationally determined densities cf. S.T. Epstein, *The Variation Method in Quantum Chemistry*, Academic Press, New York, 1974. - At degeneracies the theorem is discussed by O.E. Alon and L.S. Cederbaum, Phys. Rev. B **68**, 033105 (2003). - The Hellmann-Feynman theorem has been generalized to Gamow states by P. Ziesche, K. Kunze, B. Milek, J. Phys. A **20**, 2859 (1987).

[25] M. Isihara, Bull. Chem. Soc. Jpn. **58**, 2472 (1985).

[26] J. Vannimenus and H. F. Budd, Phys. Rev. B**15**, 5302 (1977).

[27] Z. Qian and G. Vignale, Phys. Rev. Lett. **88**, 056404 (2002); M. Polini and M. P. Tosi, Phys. Rev. B **63**, 045118 (2001); C. Caccamo, G. Pizzimenti, M. Parinello, and M. P. Tosi, Lett. Nuovo Cimento **11**, 156 (1974).

[28] J. C. Kimball, J. Phys. A **8**, 1513 (1975).

[29] H. Yasuhara and Y. Kawazoe, Physica A **85**, 416 (1976).

[30] P.-O. Löwdin, Adv. Chem. Phys. **2**, 207 (1959).

[31] M. B. Ruskai, J. Math. Phys. **11**, 3218 (1970); K. Yasuda, Phys. Rev. A **63**, 032517 (2001).

[32] P. Ziesche, in [2], p. 361; P. Ziesche, J. Tao, M. Seidl, and J. P. Perdew, Int. J. Quantum Chem. **77**, 819 (2000); P. Ziesche, in Ref. [13], p. 33. - For finite systems, particle number fluctuations in fragments have been discussed by P. Fulde, cf. Ref. [1], p. 157, further refs. are in Ref. [22].

[33] A. Rajagopal, J. C. Kimball, and M. Banerjee, Phys. Rev. B **18**, 2339 (1978).

[34] For the cumulant expansion of reduced densities and reduced density matrices cf. e.g. P. Ziesche, in [13], p. 33.

[35] P. Ziesche, Phys. Lett. A **195**, 213 (1994); Int. J. Quantum Chem. **60**, 1361 (1996); in [2], p. 361; M. Levy and P. Ziesche, J. Chem. Phys. **115**, 9110 (2001).

[36] The function $p(k) = 3k^2\mu(k)$ appears also in the mentioned fluctuation-correlation analysis of the spin-unpolarized HEG [32] and in Ref. [37].

[37] P. M. W. Gill, A. M. Lee, N. Nair, and R. D. Adamson, J. Mol. Struct. (Theochem) **506**, 303 (2002).

[38] P. Ziesche, Int. J. Quantum Chem. **90**, 342 (2002); in Ref. [3], p. 307

[39] M. Nakata, H. Nakatsuji, M. Ehara, M. Fukuda, K. Nakata, and K. Fujisawa, J. Chem. Phys. **114**, 8282 (2001); cf. also H. Nakatsuji in Ref. [13], p. 85.

[40] D. A. Mazziotti, Phys. Rev. A **65**, 062511 (2002); cf. also Ref. [13], p. 139.

Captions of the Figures

Fig.1. Dimensionless 1-matrix $f_s(r)$ of the spin-polarized HEG for different values of the spin-polarization ζ and the interaction strength r_s. r in units of $k_{\rm F}^{-1}$. Upper panel: ideal Fermi gas, $s=\uparrow$, and increasing ζ. Middle panel: ideal Fermi gas, $\zeta = 1/3$, and $s=\uparrow,\downarrow$. Lower panel: $\zeta = 1/3$, $s=\uparrow$, and increasing r_s.

Fig.2. Momentum distributions $n_s(k, r_s, \zeta)$ of the spin-polarized HEG according to Eq.(18) for the spin-polarization $\zeta = 1/3$ and the interaction-strength values $r_s = 1, 5, 10$. k in units of $k_{\rm F}$.

Fig.3. Kinetic correlation energy $t_{\rm corr}(r_s)$ in a.u. according to Perdew/Wang [5] (full line) and from $n_s(k, r_s, \zeta)$ of Eq.(18) via $t = t_\uparrow + t_\downarrow$ (the dots).

Fig.4. Non-idempotency matrices $\beta_s(r)$ and $\beta_a(r)$ of the spin-polarized HEG according to Eq.(19). r in units of $k_{\rm F}^{-1}$. 1st and 2nd panel: ideal Fermi gas, 3rd and 4th panel: increasing interaction strength r_s.



1.2
1
0.8
0.6
0.4
0.2
0
-0.2
0
2
4
6
8
10
12
14
r
$f_s(r)$
$r_s = 0$
s = ↑
ζ = 0
ζ = 1/3
ζ = 1
$r_s = 0$
ζ = 0
ζ = 1/3
↓
↑
ζ = 1/3
s = ↑
$r_s = 0$
$r_s = 5$
$r_s = 10$


1
0.8
0.6
0.4
0.2
0
0
0.5
1
1.5
2
k
$n_s(k,1,1/3)$


1
0.8
0.6
0.4
0.2
0
0
0.5
1
1.5
2
k
$n_s(k,5,1/3)$


1
0.8
0.6
0.4
0.2
0
0
0.5
1
1.5
2
k
$n_s(k,10,1/3)$



0.14
0.12
0.1
0.08
0.06
0.04
0.02
0
t_{corr} (r_s)
0
1
2
3
4
5
6
7
8
9
10
r_s



$r_s = 0$
s = ↑
ζ = 0
ζ = 1/3
ζ = 1
$\beta_s(r)$
1.2
1
0.8
0.6
0.4
0.2
0
-0.2
0
2
4
6
8
10
12
14
r



$r_s = 0$
ζ = 0
ζ = 1/3
↑
↓
$\beta_s(r)$
1.2
1
0.8
0.6
0.4
0.2
0
-0.2
0
2
4
6
8
10
12
14
r



ζ = 1/3
s = ↑
$r_s = 0$
$r_s = 5$
$r_s = 10$
$\beta_s(r)$
1.2
1
0.8
0.6
0.4
0.2
0
-0.2
0
2
4
6
8
10
12
14
r



ζ = 1/3
$r_s = 0$
$r_s = 5$
$r_s = 10$
$\beta_a(r)$
1.2
1
0.8
0.6
0.4
0.2
0
-0.2
0
2
4
6
8
10
12
14
r